

Food System Transformation in Mozambique: An Assessment of Changing Diet Quality in the context of a Rising Middle Class

By: Jenny Cairns Smart, David Tschirley and Francis Smart

Selected Paper prepared for presentation at the 2018 Agricultural & Applied Economics Association Annual Meeting, Washington, D.C., August 5-August 7

AUTHORS

Jenny Smart (j.smart@cgiar.org) was a Survey Data Analysis Advisor in the Agricultural, Food and Resource Economics Department of Michigan State University, East Lansing, when she wrote most of this work. She is currently a Research Analyst II in the Development Strategy and Governance Division of the International Food Policy Research Institute, Washington, D.C., USA.

David Tschirley (tschirle@msu.edu) is a Professor of International Development and the Co-director of the Food Security Group in the Agricultural, Food and Resource Economics Department of Michigan State University, East Lansing, MI, USA.

Francis Smart (smartfra@msu.edu) is a PhD candidate in the Measurement and Quantitative Measures division of the Counseling, Educational Psychology, and Special Education Department of Michigan State University, East Lansing, MI, USA.

Abstract

Robust income growth combined with the highest urban population growth in the world is driving very rapid changes in the food system of Sub-Saharan Africa. Demand is increasing for higher quality foods, including fresh produce, meat and dairy products as well as more processed foods, with poorer nutritional value. The overweight and obesity epidemic that first began among developed nations is not sparing the expanding middle classes within developing countries, leading to a double burden of over and under nourished populations in these areas. As rapidly expanding towns and cities proliferate across Sub-Saharan Africa, urban areas can also become deserts for fresh or less-processed nutritious foods. Urban farming has been one way that the food desert challenge in urban areas is ameliorated, and in Mozambique, even in the largest city center of Maputo, one in ten households owns their own farm land. In the context of rapid urbanization and income growth in Mozambique, this paper finds that both growing incomes and the consumption of processed foods are associated with a worsening of negative factors in the diet. Furthermore, urbanization, controlling for income, is associated more strongly with a worsening of negative factors than with an improvement in positive factors in the diet. However the effect on nutrition of owning one's own farm, controlling for the share of others in the household's area that have a farm, is positive and significant for urban households, primarily driven by these households purchasing fewer unhealthy foods. These findings have important implications concerning the role of urban farming for improving dietary quality.

Key words:

Nutrition, Dietary diversity, Dietary quality, Urbanization, Food Deserts, Urban Farming, Mozambique

Acknowledgments:

Financial and substantive support for this study was provided by the United States Agency for International Development (USAID) through the Mozambique Policy Analysis and Planning Capacity for Improved Food Security and Nutrition Outcomes (MOZCAPAN) project. The opinions expressed in this document are the authors' responsibility and do not reflect the official position of USAID.

1. Introduction, Research Questions & Hypotheses

Robust income growth combined with the highest urban population growth in the world is driving very rapid changes in the food system of Sub-Saharan Africa. Demand is increasing for higher quality foods, including fresh produce, meat and dairy products as well as more processed foods, with poorer nutritional value. The overweight and obesity epidemic that first began among developed nations is not sparing the expanding middle classes within developing countries, leading to a double burden of over and under nourished populations in these areas. As rapidly expanding towns and cities proliferate across Sub-Saharan Africa, urban areas can also become deserts for fresh or less-processed nutritious foods. Urban farming has been one way that the food desert challenge in urban areas is ameliorated, and in Mozambique - the focus of this paper - even in the largest city center of Maputo, one out of ten households owns their own farm land. In the context of rapid urbanization and income growth, this paper seeks to understand the key drivers and likely evolution of diet quality in Mozambique, in both its positive and negative dimensions, while specifically examining the potential role that urban farming might play within this transition.

Previous literature has looked at diet quality and its relationship to income, food prices, food expenditure, plot size and crop diversity among farmers, as well as a variety of household socio-demographic characteristics (Rashid et al 2011, Gaiha et al 2014, Darmon & Drewnowski 2008, Mabli et al 2010, Sodjinou et al 2009, Delisle et al 2009, Delisle et al 2010). But no one has analytically showed a relationship between own farming, city size, and nutritional outcomes. Furthermore, of those who have looked at urban farming and nutrition, no dietary quality measures such as those we developed for this study have been used (Zezza & Tasciotti 2010, Maxwell 1995, Armar-Klimesu 2000, Foeken and Mwangi 2000, Madaleno 2000, Bellows et al 2003).

We suggest five hypotheses in this study. One, the consumption of processed foods is significantly associated with worsening of negative factors in the diet. Two, income growth is associated with simultaneous improvement in positive dimensions of diet and worsening of negative dimensions. Three, the income effect on diet does not significantly differ across rural- and urban areas. Four, urbanization, (while controlling for income), is associated more strongly with a worsening of negative factors than with an improvement in positive factors. Five, not all households who are interested in eating fresh foods are able to achieve access to land to do so, however among those that are that are, having one's own farm allows the household to achieve an even more healthy diet as a result.

The rest of this paper is structured as follows: Section two describes the data and methods used in the analysis. Section three gives some descriptive statistics illustrating initial results from two methods which were explored and compared. Section four gives the output of the analysis, which is two-fold: the first section of the results will give an assessment of the patterns and drivers of current household level dietary quality (using Imamura's measure as well as others) over space (rural, large cities, secondary cities), household income levels, household education, and other demographic variables. The second section of the results anticipates the likely directions of change in diet quality over these same dimensions, based on expected income growth and expenditure elasticities developed for several alternative food groupings. By using several approaches to the projection, we are able to assess how robust the expectations can be regarding patterns of near-term evolution of diet quality. Section five summarizes our findings and concludes.

2. Data & Methods

This study uses Mozambique's nationally representative 2008/09 IOF (*Inquérito de Orçamento Familiar*) household budget survey, which was conducted from September of 2008 to September of 2009 to assess current levels, patterns, and drivers in diet quality, and to anticipate directions of change over the next

years. The IOF survey is an expenditure survey conducted in a similar style to a Living Standards Measurement Study (LSMS), and includes a large number of food items (384 unique food items in 2008/09), thus lending itself well to such analysis. These items include those purchased for household consumption, those consumed from the household's own agricultural production, and those acquired by means of compensation, for example as a payment or partial payment for work performed or meals offered to employees on the job. Responses were recorded by the household themselves over a three day period of time, or if the the respondents were not able to fill out the questionnaire themselves, the data represent a three-day recall period completed by the enumerator on their second interview visit to the household. The analysis methods we employ build on the pioneering work by Jack Fiedler and colleagues on use of expenditure surveys for dietary assessments.

We also adapt Imamura et al.'s diet quality measure for this data set (2015), which is a combined score calculated by assigning to each household a combination of positive points on the basis of increasing acquisition of nine healthy food or nutrient items (wholegrains, fruit, vegetables, fish, nuts/seeds, beans/legumes, fiber, calcium, and polyunsaturated fat) and negative points on the basis of acquisition of six unhealthy food/nutrient items (sugar-sweetened beverages, unprocessed red meat, processed meat, cholesterol, sodium and saturated fat). Corresponding applied nutritional parameters come primarily from the U.S. Department of Agriculture (USDA) food composition table, complemented when needed by other sources, including a more limited Mozambique-specific food composition table produced by Korkalo et al in 2011.

Two alternative methods are used to normalize household food quantity acquisition. In the first method, we take the approach of Smith and Subandoro (2007), among others, to generate daily per adult equivalent quantities in unit weights, of foods and nutrients acquired (whether purchased, received in kind or produced from one's own farm). In the second method, we follow the example of Mabli et al. (2010) who suggest that methods using per adult equivalent standardizations, as compared to daily recommended

values, for example, is a method that has fallen into disfavor, and recommend normalizing consumption to a 1,000 kilocalorie diet over all recorded food acquisition by the household, thus removing any adult equivalent or daily standardization. In this second approach, household size in adult equivalents is then included as a right-hand-side variable in the multi-variate analysis. In both approaches, quantities are adjusted for edible percentages and account for milliliter to gram conversion factors of liquid food items. We find the results from applying both methods to the generation of various dietary quality indicators to be useful, and where our regression analyses are carried out with both approaches, we find very similar and robust results between the two.

3. Descriptive Statistics

Using the per adult equivalent normalization approach, the first of the two methods described above, implied daily kilocalorie consumption per adult equivalent at the household level is a reasonable 2,056 kcal median, or 2,220 kcal/day mean. The six most common foods households mention acquiring within the one week recall period of the survey are vegetables (93% of all households), followed by cereals, fruit, nuts, fish, and roots (see table 3.1). Meats ranked 10th out of 13 categories, with 31% of households consuming meat during the given week.

In terms of the density of foods eaten in grams, cereal acquisition ranked highest at an average 300.5g/day/adult equivalent (primarily uncooked grains, although bread and other processed or prepared cereal products are also represented here, for example muffins or cake), followed by fruit, vegetables, roots, nuts then fish, at an average 29.4g/day/adult equivalent.

Using the same processing/perishability categorizations generated for cross-country studies such as those performed by Tschirley et al (2015a, 2015b), foods that are perishable and unprocessed are the most commonly acquired items (the group into which many fruits and vegetables are categorized), with a 70%

participation rate (see table 3.2). Perishable and highly processed foods were the least commonly consumed items, with a 31% participation rate.

Table 3.1 Most commonly represented food groups purchased among Mozambique households

Food Group	Participation Rate	Mean (g)	Median (g)	Std Dev	COV	Participation Rate Ranking
Vegetables	93%	160.8	61.5	0.54	3.38	1
Cereals	87%	300.5	322.2	0.21	0.68	2
Fruit	84%	186.0	94.1	0.38	2.06	3
Nuts	78%	61.9	36.2	0.09	1.43	4
Fish	72%	29.4	10.8	0.06	2.04	5
Roots	53%	130.1	24.3	0.24	1.84	6
Miscellaneous	41%	10.0	0.0	0.05	6.59	7
Oil	49%	7.0	0.0	0.02	2.15	8
Sugars	36%	14.8	0.0	0.04	2.45	9
Meat	31%	20.7	0.0	0.07	3.22	10
Beverages	21%	17.1	0.0	0.23	13.55	11
Egg	8%	1.2	0.0	0.01	6.27	12
Milk	3%	1.0	0.0	0.01	10.52	13

Source: 2008/2009 IOF data

Notes: Foods are categorized according to Zezza and Tasciotti's 13 comprehensive food groups (see Annex 1). Quantities are daily per adult equivalent edible percentage kgs acquired. The coefficient of variation (COV) for the categories of beverages, milk, vegetables and fruits are high due to the standardization to a 2,000 kcal diet, which in many cases causes the quantities of these high density, low calorie items, to be overrepresented in the household's collective diet.

Table 3.2 Most commonly represented processing/perishable groups purchased among Mozambique households

Processing / Perishable Food Categories	Participation Rate	Mean (g)	Median (g)	Std Dev	COV	Participation Rate Ranking
Unprocessed and Perishable	70%	76.7	34.5	0.11	1.39	1
High Processed and Non Perishable	66%	99.1	10.6	0.18	1.82	2
Low Processed and Non Perishable	61%	89.3	22.6	0.14	1.62	3
Unprocessed and Non Perishable	56%	83.3	16.5	0.14	1.68	4
Low Processed and Perishable	55%	13.7	2.3	0.03	2.47	5
High Processed and Perishable	31%	7.6	0.0	0.04	5.58	6

Source: 2008/2009 IOF data

Notes: Foods are categorized according to Tschirley et al.'s six processing and perishable categories (2015). Quantities are daily per adult equivalent edible percentage kgs acquired.

Table 3.3 Farm ownership and city size

City Size where Household is located	Someone in the household owns land	
Rural	4,301	98%
Small Town (<100k)	2,542	83%
Secondary city (100k-999k)	1,094	49%
Large city (> 1m)	152	13%
Total (N = 10,877)	8,089	74%

Source: 2008/2009 IOF data

Perhaps not surprisingly, the share of households owning land falls significantly with the household's location in a city of increasing size (see table 3.3). As we will see in the next section, household nutrition is also monotonically more negative with increasing city size. Fifty-eight percent of urban households own a farm compared to 98% in rural areas.

4. Results

4a. Patterns and Drivers of Current Household Level Dietary Quality

This section assesses the patterns and drivers of current household level dietary quality (using our adapted Imamura measure as well as others) over space (rural, small towns, secondary cities, and large cities), household income levels (proxied by total household expenditure), household education, and other demographic variables.

Indicators of Dietary Diversity

First, we briefly describe some of the key indicators of dietary diversity suggested in the literature, which we replicate or alter for our regression analysis as alternative outcome variables. Table 4.1 summarizes these indicators.

The primary measure of dietary quality we use is adapted from the measure created by Imamura et al. (2015). It is a score calculated by assigning to each household a combination of positive points on the basis of increasing acquisition of nine healthy food or nutrient items (wholegrains, fruit, vegetables, fish, nuts/seeds, beans/legumes, fiber, calcium, and polyunsaturated fat) and negative points on the basis of acquisition of six unhealthy food/nutrient items (sugar-sweetened beverages, unprocessed red meat, processed meat, cholesterol, sodium and saturated fat). The resulting value is then standardized to a 100 point scale. It is represented by the variable *imindex_std*.

We also decompose the adapted Imamura et al. combined diet quality measure into both positive and negative dimension individual scores (*imposindex_std* and *imnegindex_std*), with higher scores always indicating a better dietary outcome, whether it is increasing consumption of positive food items or decreasing consumption of negative food items. These are also standardized to a 100 point scale.

We generate and compare several dietary diversity scores, computed in one of three ways: (1) according to the number of different comprehensive food groups represented by each household (a method referenced by Sodjinou; Smith & Subandoro 2007; and Zezza & Tasciotti 2010, represented in the analysis by variables *discr13* and *divscr7*), (2) as a simple count of the different food items the household reported having acquired (a method employed by Zezza & Tasciotti 2010, and represented by the variable *foodcount*), or (3) via several alternative versions of a food diversity index (variables *fdi13quant*, *fdi13expen*, *fdi7quant*, and *fdi10quant*), which are calculated as the sum of squares of the shares of the various food items in the food consumption basket (Gaiha et al 2012). These are created separately as a function of food quantities and as a function of food expenditures. The regression results in relation to the different dietary diversity score specifications are robust in that they yield quite similar estimates.

Table 4.1 Regression Outcome Variables

Dietary Quality Indicators	Definition/Further Explanation	Obs	Mean	Std. Dev.	Min	Max
Imamura Indices	Combination scores of positive points for positive items consumed, and negative for negative items consumed. Scores of 1-5 given on the basis of the food's per adult equivalent consumption quintile.					
<i>imindex_std</i>	Imamura Combined Index – a combined score of the consumption of healthy and unhealthy foods.					
<i>imposindex_std</i>	Imamura Positive Index (higher score = more consumption of healthy foods).					
<i>imnegindex_std</i>	Imamura Negative Index (higher score = less consumption of unhealthy foods).					
Dietary Diversity Scores	Defined as the total number of different food groups consumed over a given recall period, irrespective of the frequency and the amounts consumed.					
<i>divscr13</i>	Calculated on the basis of the food categorizations taken from Zezza and Tasciotti (2010), this is an exhaustive categorization of 13 subcomponents including sweets as one and oils/fats as another.	10876	6.6	2.1	1.0	13.0
<i>divscr7</i>	Similar to Gaiha et al's five groups (2012), this indicator excludes any beverages or misc group, and is defined over the following seven food groups, as offered by Smith and Subandoro (2007): 1 - cereals, roots and tubers, 2 - pulses and legumes, 3 - dairy products, 4 - meats, fish and seafood, and eggs, 5 - oils and fats, 6 - fruits, and 7 - vegetables.	10876	4.5	1.2	0.0	7.0
<i>foodcount</i>	A simple count of the different food items the HH reported having consumed during the reference period.	10876	11.0	4.8	1.0	43.0
Food Diversity Indices	Taken from Gaiha et al, this is constructed as the sum of squares of the shares of the various food items in the food consumption basket. A high value of the index implies a more concentrated food basket and low value implies a more diverse food basket. The five food groups to construct the FDI in Gaiha et al are: (i) cereals and pulses; (ii) milk, milk products, eggs, and meats; (iii) oil; (iv) sugar; and (v) fruits and vegetables.					
<i>fdi13quant</i>	Generated on the basis of food quantities acquired among the Zezza and Tasciotti's exhaustive categorization of 13 subcomponents (2010), including sweets as one and oils/fats as another (see annex table 6.1).	10876	0.3	0.1	0.1	1.0
<i>fdi13expen</i>	Generated on the basis of food expenditures acquired among the Zezza and Tasciotti's exhaustive categorization of 13 subcomponents (2010), including sweets as one and oils/fats as another.	10876	0.4	0.2	0.1	1.0
<i>fdi7quant</i>	Generated on the basis of food quantities acquired among the seven food groups offered by Smith and Subandoro (2007), and excludes any beverages or misc group.	10876	0.4	0.1	0.1	1.0

Dietary Quality Indicators		Definition/Further Explanation	Obs	Mean	Std. Dev.	Min	Max
<i>fdi10quant</i>		Generated on the basis of food quantities acquired among the 10 food items identified by Imamura et al, 2015: Wholegrain, fruit, vegetable, fish, nuts&seeds, beans&legumes, milk, sugar-sweetened beverages, unprocessed red meats, and processed meats (excluding fruit juice).	10876	0.4	0.2	0.1	1.0
General/Overall Indicators							
<i>AMDS</i>		Alternative Mediterranean Diet Score (AMDS or AMED), generated on the basis of the table referenced in annex table 6.2.	10876	3.3	1.3	0.0	8.0
<i>healthflsc</i>		Healthfulness Score: How consumed diets compare to the WHO/FAO recommendations in the areas of total fat, saturated fatty acids, polyunsaturated fatty acids, cholesterol, sugar, protein, fibre, fruits, vegetables, and nuts. Created on the basis of percentage contributions, for more detail, see annex tables 6.3 through 6.7.	10876	2.9	1.0	0.0	7.0
Macronutrient Diet Composition Scores							
<i>macroratiosc</i>		Macronutrient ratio score on the basis of widely accepted general guidelines as the desirable ranges of carbohydrate: protein: fat proportions. For more detail, see annex table 6.8.					
<i>monosatratio</i>		Ratio of monounsaturated to saturated fatty acids.	10876	1.7	1.6	0.0	10.8
<i>sfatpct1</i>		Saturated fat share of total energy.	10876	0.1	0.1	0.0	0.8
<i>pfatpct1</i>		Polyunsaturated fat share of total energy.	10876	0.0	0.0	0.0	0.2
<i>lippct1</i>		Total fat share of total energy.	10876	0.2	0.1	0.0	0.9

Source: 2008/2009 IOF data

Table 4.2 Determinants of Dietary Quality

VARIABLES	lnimindex	lnimposindex	lnimnegindex	divscr13	AMDS	healthflsc	macroratiosc	monosatratio
hgender	0.0297***	0.0231*	0.0367***	-0.214	0.017	0.346***	0.0252	-0.0602
hage	-9.09E-06	-7.92e-05***	6.97e-05***	-0.00196***	-0.00058***	-0.000402*	-0.000369	-0.00136***
1.citysize	-0.00728**	0.00937	-0.0229***	0.548***	0.0381	-0.0702*	0.195	0.00049
2.citysize	-0.0180***	0.0131	-0.0473***	1.041***	-0.120*	-0.184***	0.205	-0.423***
3.citysize	-0.0374***	-0.0242	-0.0529***	1.574***	-0.0338	0.190*	0.450*	-1.053***
shareownfarm	-0.0352***	-0.0262	-0.0494***	0.376	-0.322**	-0.451***	0.168	0.184
ownfarmShare ownfarm	0.0400***	-0.0154	0.103***	-0.622***	-0.14	0.268***	0.0422	-0.296**
totexp	-0.00244***	0.000248	-0.00534***	0.0401***	0.000281	0.0195***	0.0152**	0.00482
fhheduc	0.000171	0.000561	4.66E-05	0.0592***	0.0112	0.00733	-0.0105	-0.0155
mhheduc	-0.00254*	-0.00287	-0.00244*	0.0432*	-0.0147	-0.0154	0.0251	0.0374***
fsalaried	-0.0375***	-0.0386***	-0.0407***	-0.392**	-0.341***	-0.453***	-0.0552	-0.417***
fcontp	-0.0127***	-0.00141	-0.0241***	0.291***	0.0406	-0.0810*	-0.0923	-0.154***
fnowork	-0.0201***	-0.0179*	-0.0235***	-0.0692	-0.152**	-0.279***	0.0265	-0.178**
msalaried	0.00629	0.022	-0.00955	0.292*	0.154	0.0599	0.143	-0.0634
mcontp	0.0188***	0.0243**	0.0155*	0.420***	0.184**	0.358***	-0.0765	-0.0627
mnwork	0.0199*	0.0203	0.0205	-0.127	0.0864	0.163	-0.199	-0.333***
nkidslt5	-0.00450***	-0.00517*	-0.00386**	0.116***	-0.000271	-0.0438**	0.0451	0.0107
nkids1220	-0.00504***	-0.00579*	-0.00396**	0.109***	-0.0247	-0.0840***	-0.0347	-0.0677***
nummeals	0.00397	0.0263***	-0.0176***	0.571***	0.208***	0.157***	-0.0502	0.0713***
Constant	3.867***	3.113***	3.213***	3.561***	2.935***	3.297***	2.789***	1.932***
R-squared	0.061	0.014	0.185	0.174	0.02	0.042	0.025	0.042

Significance at 1% (***); 5% (**); 10% (*).

The Alternative Mediterranean Diet Score (*AMDS*) and the Healthfulness Score (*healthflsc*) are two overarching scores which assign a value based on how consumed diets compare to the WHO/FAO recommendations in the areas of total fat, saturated fatty acids, polyunsaturated fatty acids, cholesterol, sugar, protein, fiber, fruits, vegetables, and nuts (Sodjinou et al 2009, Deslise et al 2009). The AMDS is fashioned after Fung et al.'s application in their article from the journal of clinical nutrition (2005).

The Macronutrient Ratio Score (*macroratiosc*) is calculated on the basis of conformance to the widely accepted general guidelines of desirable ranges for energy proportions from carbohydrates, protein and fat (Kim et al 2003, Deslise et al 2009). Lastly, the ratio of monounsaturated to saturated fatty acids (*monosatratio*) is also used as its own indicator of dietary quality (Deslise et al 2009), a higher score being more desirable. Variables *lippct*, *sfatpct*, and *pfatpct* represent the total, saturated, and polyunsaturated fat shares of total energy.

Regression Results

Here, we summarize some of our main regression take-aways. The dependent variable definitions and full regression results are reported in annex tables 6.9 through 6.12, but a shortened set of regression results can be found in table 4.2. Note that food share-specific regression results are based on the all-inclusive food categories as suggested in Zezza & Tasciotti (2010).

Households having a female head has almost an invariably positive effect on nutrition, across diet quality or dietary diversity indicators and significant particularly in the case of the Imamura scores, both more positive and less negative dimensions, as well as the “healthfulness score” as fashioned after Sodjinou et al. Food share-specific regressions indicate that female headed households consume significantly more cereals and vegetables as well as a significantly less sugars, syrups, sweet, oils and fats.

Increasing age overall has a negative effect on nutrition, although is related to significant decreases in acquisition of both positive and negative items of the Imamura index. Increasing age has a strong and robust negative effect on dietary diversity, Mediterranean diet score, healthfulness score, and monosaturated to saturated fat ratio.

Living in an urban area has an increasingly negative effect on dietary quality the larger the city (robust across Imamura index, healthfulness score, Mediterranean diet score, and the mono/saturated fat ratio), primarily via increased negative food item consumption, but also through a decrease in positive food items. Food share-specific regressions indicate an increase specifically in sugar, fat and (non-specific) beverage shares of food expenditures as city size increases, and decreases in vegetable and nuts/seeds/legume shares. Living in an urban area has an increasingly positive effect on dietary diversity as city size increases. Food-specific regressions indicate an increase specifically in fruit shares of food expenditures in the cities over 1 million, and increasing shares of fish expenditures as city size increases from rural to small towns (less than 100k) and medium size urban centers (100-999k).

The share of those owning their own farms has a negative effect on dietary quality (both Imamura index, the healthfulness score, and the alternative Mediterranean diet score), perhaps picking up in part on farming households located in lower income urban areas via the increase in acquisition of negative items rich in oil and fats.

Owning one's own farm, controlling for the share of those who own their own farm in the surrounding area (area defined by a geographical unit with 6 to 84 households, or an average 15), has a positive and significant effect on dietary quality (both the Imamura index, the healthfulness score and a decrease in the percentage of total fat calories in the diet) via a strong and significant decrease in negative items acquired. At the same time, owning one's own farm also has a negative and significant effect on dietary diversity. This may be due to farmers relying more heavily on their own production and not visiting a

market as frequently. Food-specific regressions indicate an increase specifically in cereals/grains, starchy/roots/tubers/legumes and nut/seed/legume group shares of food expenditures and a decrease in sugar, fat, beverages and meat shares among households owning their own farms.

Total expenditure has a negative effect on dietary quality; however has mixed effects on dietary diversity between both of our methodological approaches - the per adult equivalent and 1000 calorie normalization techniques. Total expenditure has a positive effect on the healthfulness score, the macronutrient ratio score, and the poly and monounsaturated fat share of total energy measures, so these results are a bit hard to discern, and discussed further in the next section on income projections. Food group share analysis indicates an increase specifically in meat, beverages, eggs and milk product food expenditure shares and a decrease in vegetable, fish, nut/seed/legume and roots/tubers shares as well as sugary foods, as total expenditure increases.

Both male and female household head education is has a positive effect on dietary diversity, educated female heads, more strongly so. Food-specific regressions indicate an increase specifically in milk and meat expenditure shares among these households. Education of male heads specifically has a negative effect on dietary quality as it regards significant increased consumption of negative items, while also having a positive effect on the monosaturated to saturated fat ratio via a significant reduction in consumption of saturated fats in these households. The effect on these indicators of female household head education were insignificant. Food-specific regressions indicate a decrease specifically in nuts/legumes and vegetables expenditure shares and an increase in oil/fat and beverage shares among these households headed by educated males.

The number of adult equivalents in each home also has a negative effect on dietary quality as it relates to significant increased consumption of negative items as well as a lower monosaturated to saturated fat

ratio. Food-specific regressions indicate an increase in the share of sugar, syrups and sweets expenditures with increasing household size.

Male *conta propria* self-employment is the only category of employment among male or female household heads which has a significant and positive effect on nutrition quality and also shows a significant strong relationship to increased dietary diversity, an increased healthfulness and AMDS score, and a decreased saturated fat percentage of total energy. It appears these positive effects are primarily via the impact of both a significant increase in positive food or nutrients acquired and significant decrease in negative food or nutrients acquired among these households. Food group share-specific regressions indicate an increase in cereals and grain expenditures and a decrease in oil and fat expenditures among this group.

Female salaried employment is the only employment category among male or female household heads which has a significant and negative effect on nutrition quality, dietary diversity and other nutrition indicators (AMDS, healthfulness score, and mono-saturated fat ratio, via significant increases in saturated fat). Other employment categories had more mixed results, sometimes positive and sometimes negative depending on the indicator. Food group share-specific regressions indicate a significant increase in sugar and fat food group share expenditures among female household head salaried employees, however also significant increases in fish and milk shares.

Having kids less than five years old and kids ages 12 to 20 seems to have a generally negative effect on household nutrition, although this effect is not significant across all results, the strongest driver deriving from increased negative food item consumption, especially among families with adolescents.

4b. Expected Evolution of Dietary Quality as Incomes Rise

Again, using the processing/perishability categorizations mentioned in section three, now including two categories that capture own production and level of perishability, we look at the nutritional heterogeneity across these. The eight comprehensive food groupings are as follows: 1. Own produced and perishable, 2. Own produced and unperishable, 3. Unprocessed and perishable, 4. Unprocessed and nonperishable, 5. Low processed and perishable, 6. Low processed and nonperishable, 7. High processed and perishable, and 8. High processed and non-perishable. In addition to this, we also compare these eight processing/perishability categories to the three food classification groups offered by Monteiro et al according to the degree and purpose of industrial processing, listed in annex table 6.13 (2010).

Through ANOVA regressions (coefficients reported in table 4.3) we find that these eight categories do indeed explain a fair part of the Imamura diet quality index, with an R-squared of 0.27 at the community level, and of 0.20 at the household level, and all but one category significant in both. Monteiro's three groups, on the other hand, explained very little (at the household level, an R-squared of 0.12 and at the community level, an R-squared of 0.13), despite each included category being significant, which is an interesting result, since nutrition was a key motivation for Monteiro et al's classification.

Table 4.3 How much of the Imamura combined index is explained by the eight-group processing/perishability food categorization budget shares

Food Category	Level of Significance	
	HH <i>R2=0.20</i>	Community <i>R2=0.27</i>
These categories increase diet quality:		
2 Own production and non-perishable	2.3***	2.0
4 Unprocessed and non-perishable	8.2***	20.0***
These categories decrease diet quality:		
3 Unprocessed and perishable	0.4	-12.1***
5 Low processed and perishable	-8.9***	-19.6***
6 Low processed and non-perishable	-4.1***	-5.6**
7 High processed and perishable	-8.7***	-9.1***
8 High processed and non-perishable	-9.2***	-10.0***

Source: 2008/2009 IOF data

Notes: Results are relative to the "own production and perishable" default group number one. (***) $p < 0.01$, (**) $p < 0.05$, (*) $p < 0.1$

Relative to the own production and perishable default group, the only two food categorization shares which explained an improvement in the Imamura combined diet quality index from the eight described above are share two, own production and non-perishable foods, as well as share four, unprocessed and non-perishable foods. Consumption of foods in any of the low or high processed food categories corresponds to a negative effect on dietary quality (frequent items in these groups include common quality rice, white corn maize, normal wheat bread, traditional liquor, wine, etc).

How is dietary quality then likely to evolve given expected income growth?

By looking at expenditure elasticities developed for several alternative food groupings, we begin to address the question of how dietary quality is likely to evolve given expected income growth.

While the results on the overall expenditure relationship to dietary quality are mixed in the analysis of the drivers and patterns of dietary decisions above, the food expenditure elasticity to the Imamura et al dietary quality indices, by comparison, are positive and insignificant. That is to say, for a 10% increase in food spending, the nutritional value of food purchased as measured by this index is ambivalently better. However for an increase in overall spending (a proxy for income level), dietary quality, as measured by the Imamura indices, decreases.

The elasticity of demand for several essential vitamins and minerals, fiber, and other micronutrients, show that several, if not most, of these items are inferior goods. While vitamin A is projected to (dramatically) increase with increasing expenditure on food, vitamins C, Riboflavin and Vitamin B12 are projected to substantially decrease, and these appear to be much lower than the trajectories of the lowest-income group of Americans in Mabli et al's study, using data from 1996. A full list of vitamin and mineral elasticities

given a 1% increase in total daily expenditure in Purchase Power Parity per adult equivalent on food is given in table 4.4.

There can be no doubt that the dropping vegetable, as well as fruit food shares – although the latter, insignificantly so, which are being driven by increasing total household expenditures are also related to the negative elasticities for many of these essential vitamins and minerals. And as mentioned in the previous section, the food group expenditure shares of roots/tubers/legumes, nuts/seeds, cereals/grains and fish/shellfish are also expected to decrease with increasing total expenditure, while meat, egg and beverage shares increase.

Table 4.4 Vitamin and Mineral Food Expenditure Elasticities

	N	% missing values	Food Expenditure Elasticity
Vitamin A iu	10,836	0.4%	0.38
Vitamin B6 mg	10,872	0.0%	0.05
Vitamin B12 g	8,751	19.5%	-0.30
Vitamin E mg	10,855	0.2%	-0.09
Total Folate g	10,869	0.1%	-0.08
Niacin mg	10,872	0.0%	-0.01
Riboflavin (B2) mg	10,872	0.0%	-0.50
Vitamin C mg	10,813	0.6%	-0.44
Calcium mg	10,875	0.0%	-0.11
Iron mg	10,874	0.0%	-0.19
Magnesium mg	10,872	0.0%	-0.02
Phosphorous mg	10,873	0.0%	0.01
Zinc mg	10,871	0.0%	0.08
Sodium mg	10,875	0.0%	-0.11

Note: All significant at the $p < .01$ level except for Niacin and Phosphorous.

Source: 2008/2009 IOF data

We can see from these results, that as incomes rise and Mozambique's expenditure capacity increases, there will be significant and positive increases in the shares of purchased food which is highly processed, and are shown to decrease dietary quality. This will be concurrently accompanied by a significant drop in

shares of own production of food consumed, as well as a drop in the share of food purchased which is unprocessed and nonperishable, both of which are associated with significantly increasing dietary quality.

5. Conclusion

In conclusion, we return to our original five hypotheses. One, as we hypothesized, the consumption of processed foods is significantly and strongly associated with the worsening of negative factors in the diet.

Two, income growth is associated with the worsening of negative dimensions of diet, however is not consistently associated with simultaneous improvement in positive dimensions of diet, nor are these results, when positive, significant. In fact, some of our results do show the troubling result of significant decreases in consumption of positive dimensions of diet with households' rising income.

Three, whether the income effect on diet does significantly differ across rural and urban areas varies based on the method of standardization we applied to household food acquisition. When the approach is used that normalizes total food acquisition to a 1,000 calorie diet, the income effect on diet is negative and significant and this does not significantly differ across rural and urban areas, as we hypothesized. However when the approach is used that normalizes food acquisition by day and per adult equivalent, the income effect is negative and significant in the urban areas, but positive and significant in the rural areas.

Four, as we hypothesized, urbanization, controlling for income, is associated more strongly with a worsening of negative factors than with an improvement in positive factors.

Five, we assumed that households interested in eating fresh foods are more likely to seek out land to farm so that they can eat more fresh food, regardless of if they are actually able to achieve access to land to do so or not. We then asked, for those that do gain access to farm land, does this allow them to eat even

more healthy diets than they would otherwise? Our results show that as we hypothesized, controlling for the share of others in one household's area that have a farm, the effect on nutrition of owning one's own farm is positive and significant, primarily driven by these households purchasing lesser amounts of negative food items.

As the Mozambican middle class grows and urbanization continues, it will be an uphill battle for households to prioritize good nutritional choices in their diets, and this analysis shows that particularly vulnerable groups may be educated and/or salaried females and parents of adolescents and children under five. Especially in urban settings, sugar and fat dense foods may very well become the most accessible and convenient food to be found, however, are lacking in essential nutrients. Yet even in urban contexts, there are many households who have been able to find ways of growing some of the food they consume for themselves, and this study shows that doing so is associated with these same households acquiring more healthy items in their diets in general, and achieving even greater nutrition than their counterparts. Further research is needed on the topic of what opportunities urban farming may present, moving into the future, to prevent highly processed and unhealthy foods from crowding out households' opportunities to consume a nutritious whole foods diet.

6. Annex Tables

The following tables provide supplementary material for the text of the main body of this paper.

Annex Table 6.1. Zezza and Tasciotti's 13 comprehensive food groups (2010)

Food group	Content of the group
Food group 1	Cereals and grain products
Food group 2	Starchy, roots, tubers and legumes
Food group 3	Nuts, seeds and legumes
Food group 4	Vegetables
Food group 5	Fruits
Food group 6	Sugar, syrup and sweets
Food group 7	Meat and poultry
Food group 8	Fish and shellfish
Food group 9	Milk and milk products
Food group 10	Oil and fats
Food group 11	Beverages
Food group 12	Eggs
Food group 13	Miscellaneous

Annex Table 6.2. The Alternate Mediterranean Diet Score, as defined by Fung et al. (2005)

Food Group	Foods Included	Criteria for 1 point ^a
Vegetables	All vegetables except potatoes	Greater than median intake (Servings/d)
Legumes	Tofu, string beans, peas, beans	Greater than median intake (Servings/d)
Fruit	All fruit and juices	Greater than median intake (Servings/d)
Nuts	Nuts, peanut butter	Greater than median intake (Servings/d)
Whole grains	Whole-grain ready-to-eat cereals, cooked cereals, crackers, dark breads, brown rice, other grains, wheat germ, bran, popcorn	Greater than median intake (Servings/d)
Red and processed meats	Hot dogs, deli meat, bacon, hamburger, beef	Greater than median intake (Servings/d)
Fish	Fish and shrimp, breaded fish	Greater than median intake (Servings/d)
Ratio of monounsaturated to saturated fat	-	Greater than median intake (Servings/d)
Ethanol	Wine, beer, "light" beer, liquor	5-25 g/d

^a 0 points if these criteria are not met.

How the Healthfulness Dietary Quality Score was generated relative to FAO/WHO guidelines

Annex Table 6.3 The FAO/WHO guideline ranges of population nutrient intake goals (2003)

Dietary factor	Goal (% of total energy, unless otherwise stated)
Total fat	15-30%
Saturated fatty acids	< 10%
Polyunsaturated fatty acids (PUFAS)	6-10%
n-6 Polyunsaturated fatty acids (PUFAS)	5-8%
n-3 Polyunsaturated fatty acids (PUFAS)	1-2%
Trans fatty acids	<1%
Monounsaturated fatty acids (MUFAS)	By difference ^a
Total carbohydrate	55-75% ^b
Free sugars ^c	< 10%
Protein	10-15% ^d
Cholesterol	<300 mg per day
Sodium chloride (sodium) ^e	<5g per day (<2g per day)
Fruits and vegetables	> 400 g per day
Total dietary fibre	From foods ^f
Non-starch polysaccharides (NSP)	From foods ^f

^a This is calculated as: total fat – (saturated fatty acids + polyunsaturated fatty acids + trans fatty acids)

^b The percentage of total energy available after taking into account that consumed as protein and fat, hence the wide range.

^c The term “Free sugars” refers to all monosaccharides and disaccharides added to foods by the manufacturer, cook or consumer, plus sugars naturally present in honey, syrups and fruit juices.

^d The suggested range should be seen in the light of the Joint WHO/FAO/UNU Expert Consultation on Protein and Amino Acid Requirements in Human Nutrition, held in Geneva from 9 to 16 April 2002.

^e Salt should be iodized appropriately. The need to adjust salt iodization, depending on observed sodium intake and surveillance of iodine status of the population, should be recognized.

^f See page 58 of the full report, under “Non-starch polysaccharides”

Sodjinou et al, in their 2009 study, create a healthfulness score by assigning a score of 1 for each item if the recommendation was met, and 0 if it was not, for a maximum of 8. Similarly, in this study we give a score of 1 for each item if the recommendation was met, per the following list, also for a maximum of 8.

Annex Table 6.4 Sodjinou et al’s criteria for generating their “Healthfulness Score”

Fat percent of total energy falls between 15 and 30%
Saturated fat percent of total energy is lower than 10%
Polyunsaturated fat percentage of total energy is between 6-10%
Carbohydrate percentage of total energy is between 55 and 75%
Protein percentage of total energy is between 10 and 15%
Cholesterol acquisition is lower than 300 mg/day
Fruit and vegetable acquisition is greater than 400mg/day
Fiber acquisition is greater than 25g/day

Annex Table 6.5 Our Healthfulness Dietary Quality Score Individual component results

Variable	Obs	Mean	Std. Dev.	Min	Max
Fat percent of total energy	10745	0.23	0.09	0.00	0.85
Saturated fat percent of total energy	10745	0.07	0.06	0.00	0.76
Polyunsaturated fat percentage of total energy	10745	0.04	0.02	0.00	0.19
Carbohydrate percentage of total energy	10745	0.64	0.11	0.00	1.00
Protein percentage of total energy	10745	0.23	0.08	0.00	0.81
Daily Cholesterol acquisition	10745	509.24	858.11	0.00	29086.58
Fruit and Vegetable acquisition	10745	328.14	330.99	0.00	3402.35
Daily fiber acquisition	10745	40.55	29.96	0.00	253.42

Annex Table 6.6 Our Healthfulness Dietary Quality Score Distribution

Healthfulness Score	Frequency	Percent	Cumulative Percent
0	66.00	0.61	0.61
1	356.00	3.31	3.93
2	1334.00	12.42	16.34
3	2701.00	25.14	41.48
4	3382.00	31.48	72.95
5	2242.00	20.87	93.82
6	607.00	5.65	99.47
7	57.00	0.53	100.00

Note that this method is also similar to the criteria used for the healthy diet indicator offered by Huijbregts et al. (1997), based on the dietary guidelines for the prevention of chronic diseases.

Annex Table 6.7 The Healthy Diet Indicator offered by Huijbregts et al.

Nutrient or food group (daily intake)	Dichotomous value	
	1	0
Saturated fatty acids	0-10	>10
Polyunsaturated fatty acids	3-7	<3 or >7
Protein	10-15	<10 or >15
Complex carbohydrates	50-70	<50 or >70
Dietary fibre (g)	27-40	<27 or >40
Fruits and vegetables (g)	>400	<400
Pulses, nuts, seeds (g)	>30	<30
Monosaccharides and disaccharides	0-10	>10
Cholesterol (mg)	0-300	>300

Note: Values are percentage of energy intake unless indicated otherwise.

Annex Table 6.8. Macronutrient ratio score cutoff values, per Kim et al, 2003

Macronutrient ratio (carbohydrate: protein: fat), for 0 to 6 points:

55 ~ 65: 10 ~ 15: 15 ~ 25 = 6

52 ~ 68: 9 ~ 16: 13 ~ 27 = 4

50 ~ 70: 8 ~ 17: 12 ~ 30 = 2

Otherwise = 0

Widely accepted general guidelines were chosen as the desirable ranges for the proportions of energy from carbohydrates, protein and fat.

Annex Table 6.9. Independent Variables used in the Regression Analysis

hgender	HH head gender (0 = male, 1 = female)
hage	HH head age
citysize	0 = rural, 1=small town (<100k), 2=secondary city (100k – 1m), 3=large city (> 1m)
ownfarm-smallcity	City size interacted with variable for if anyone in the HH owned his or her own farm
ownfarm-medcity	City size interacted with variable for if anyone in the HH owned his or her own farm
ownfarmlgcity	City size interacted with variable for if anyone in the HH owned his or her own farm
shareownfarm	Share of HHs in district-aldeia/bairro-urban area with at least 1 family member owning farmland
ownfarmShare-ownfarm	" <i>Shareownfarm</i> " interacted with <i>ownfarm</i> dummy variable for if anyone in the HH owned his or her own farm
totexp	Total daily expenditure in per adult equivalent
fhheduc	Highest level of education completed by female head or female spouse of head
mhheduc	Highest level of education completed by male head or male spouse of head
fsalaried	Female head or female spouse of head works for govt, public or private sector (dummy, 1 = yes 0 = no)
fcontap	Female head or female spouse of head works <i>conta própria</i> (self employed) with or without employees (dummy, 1 = yes 0 = no)
fnowork	Female head or female spouse of head does not work (dummy, 1 = yes 0 = no)
msalaried	Male head or male spouse of head works for government, public or private sector (dummy, 1 = yes 0 = no)
mcontap	Male head or male spouse of head works <i>conta própria</i> (self employed) with or without employees (dummy, 1 = yes 0 = no)
mnowork	Male head or male spouse of head does not work (dummy, 1 = yes 0 = no)
nkidslt5	Number of kids under age 5
nkids1220	Number of kids between 12 and 20
nummeals	Number of meals the family shared the previous day (0-3)

Annex Table 6.10 Full Regression Results

VARIABLE S	lnimindex	lnimpoindex	lnimnegindex	divscr13	divscr7	foodcount	fdi13q	fdi7qu	fdi13e	fdi10q	AMDS	healthfisc	macro ratios	monosatratio	sfatpct1	pfatpct1	lippct1
hgender	0.0297***	0.0231*	0.0367***	-0.214	0.0301	-0.194	-0.00659	-0.0182	0.0173	-0.0217*	0.017	0.346***	0.0252	-0.0602	-0.00775	-	0.0161**
hage	-9.09E-06	-7.92e-05***	6.97e-05***	0.00196**	0.000873**	0.00217**	0.000135***	0.000153***	0.000130***	0.000110***	0.000582**	0.000402*	0.00036*	0.00136**	-5.51E-06	-1.96e-05***	-9.43e-05***
1.citysize	-0.00728**	0.00937	0.0229***	0.548***	0.317**	0.716***	0.0181**	0.0145**	0.0239**	0.0136**	0.0381	-0.0702*	0.195	0.00049	0.0102**	0.000375	0.0189**
2.citysize	-0.0180***	0.0131	0.0473***	1.041***	0.715**	2.091***	0.0385**	0.0368**	0.0687**	0.0291**	-0.120*	0.184***	0.205	-0.423***	0.0184**	-0.00054	0.0273**
3.citysize	-0.0374***	-0.0242	0.0529***	1.574***	0.761*	5.130***	0.0609**	0.0688**	0.102***	0.0492**	-0.0338	0.190*	0.450*	-1.053***	0.0206**	0.00927*	-0.00288
shareownfarm	-0.0352***	-0.0262	0.0494***	0.376	0.542**	-0.00391	0.0126	0.0257	-0.0153	0.0303	-0.322**	0.451***	0.168	0.184	0.00538	0.00404*	0.0127
OwnfarmShareownfarm	0.0400***	-0.0154	0.103***	-0.622***	-0.123	1.093***	0.0115	-0.002	0.0734**	0.0349**	-0.14	0.268***	0.0422	-0.296**	0.0037	0.00737*	0.0267**
totexp	-0.00244**	0.000248	0.00534**	0.0401**	0.0115**	0.0694**	0.00199**	0.00164*	0.00160*	0.00114*	0.000281	0.0195**	0.0152*	0.00482	6.35E-05	0.000249**	0.000105
fhheduc	0.000171	0.000561	4.66E-05	0.0592**	0.0312**	0.132***	0.00322*	0.00338*	0.00731*	0.00389*	0.0112	0.00733	-0.0105	-0.0155	0.000202	0.000184	0.000572
mhheduc	-0.00254*	-0.00287	-0.00244*	0.0432*	0.0291**	0.0786*	0.000988	0.000275	-0.00149	-0.0017	-0.0147	-0.0154	0.0251	0.0374**	0.00123*	0.000161	0.000886
fsalaried	-0.0375***	0.0386**	0.0407***	-0.392**	0.250**	-0.851**	0.0203*	0.0153	-0.00251	0.0351**	-0.341***	0.453***	-0.0552	-0.417***	0.0137**	-0.00102	0.00382
fcontp	-0.0127***	-0.00141	0.0241***	0.291***	0.175**	0.710***	-0.00765	0.00969*	0.0260**	-0.00442	0.0406	-0.0810*	-0.0923	-0.154***	0.00967**	0.00244**	0.0254**
fnowork	-0.0201***	-0.0179*	0.0235***	-0.0692	-0.0135	-0.147	0.00531	0.000596	-0.00136	0.0118	-0.152**	0.279***	0.0265	-0.178**	0.0106**	0.00129*	0.0184**
msalaried	0.00629	0.022	-0.00955	0.292*	0.259**	0.791***	0.0326**	0.0431**	-0.0196	0.0408**	0.154	0.0599	0.143	-0.0634	-0.0043	0.00166	0.000531
mcontp	0.0188***	0.0243**	0.0155*	0.420***	0.408**	0.838***	0.0324**	0.0446**	-0.00995	0.0467**	0.184**	0.358***	-0.0765	-0.0627	0.0128**	0.00219*	-0.00397
mnwork	0.0199*	0.0203	0.0205	-0.127	-0.121	-0.103	-0.0142	-0.0171	0.00133	-0.022	0.0864	0.163	-0.199	-0.333***	-0.00474	-0.00016	0.0168**
nkids15	0.00450**	-0.00517*	0.00386**	0.116***	0.0464***	0.237***	0.00706**	0.00857**	0.00546*	0.00800**	-0.000271	0.0438**	0.0451	0.0107	0.00012	0.000182	0.000959
nkids1220	0.00504**	-0.00579*	0.00396**	0.109***	0.0452***	0.333***	9.99E-05	0.00101	0.00117	0.000916	-0.0247	0.0840**	-0.0347	0.0677**	0.00269**	0.00019	0.00332**
nummeals	0.00397	0.0263**	0.0176***	0.571***	0.301*	1.300***	0.0268**	0.0260**	0.0213**	0.0345**	0.208***	0.157***	-0.0502	0.0713**	0.00478**	0.00147**	-0.00159
Constant	3.867***	3.113***	3.213***	3.561***	2.368**	4.570***	0.421***	0.506***	0.443***	0.532***	2.935***	3.297***	2.789**	1.932***	0.0746**	0.0428**	0.222***
Observations	10,873	10,873	10,873	10,873	10,873	10,873	10,873	10,873	10,873	10,873	10,873	10,873	1,783	10,873	10,873	10,873	10,873
R-squared	0.061	0.014	0.185	0.174	0.101	0.236	0.075	0.063	0.11	0.065	0.02	0.042	0.025	0.042	0.037	0.061	0.051

*** p<0.01, ** p<0.05, * p<0.1; green highlighting is a significantly positive effect, orange highlighting is a significantly negative effect

Annex Table 6.11 Full Regression Results using alternative standardization method

VARIABLES	lnimind ex	lnimposin dex	lnimnegin dex	divscr1 3	divscr7	foodco unt	fdi13qu ant	fdi7qua nt	fdi13ex pen	fdi10qu ant	monosatr atio	sfatpct l	pfatpct l	monofatsh are	lippct1			
hgender	0.0266** *	0.0269**	0.0280***	-0.168	0.039	-0.0593	-0.00996	-0.0211*	0.0222*	-0.0297**	-0.0602	-0.00775	-0.000715	-0.306	-0.0161**			
hage	-6.00E-06	-9.53e-05***	8.75e-05***	-	-	-	9.77e-05***	0.000113 ***	0.000113* **	9.09e-05***	-0.00136***	-5.51E-06	-1.96e-05***	-0.00320***	-9.43e-05***			
1.citysize	-	-	-	0.0176** *	-0.00137	-0.0314***	0.554***	0.319***	0.727***	-	-	-	-	-	-	0.0189** *		
2.citysize	0.0463** *	-0.0524***	-0.0384***	1.016***	0.707***	2.038***	-	-	-	-	-	-	-	-	-	0.0273** *		
3.citysize	0.0685** *	-0.0890***	-0.0483***	1.690***	0.812***	5.333***	0.0758***	0.0721** *	-0.0981***	0.0684***	-1.053***	0.0206* **	-	-	-	0.00927* **		
shareownfarm	0.0659** *	-0.133***	0.00337	0.365	0.523***	0.0263	0.0374**	0.0558** *	-0.0123	0.0433**	0.184	0.00538	-	-	-	0.00404* *		
OwnfarmShare ownfarm	0.0511** *	0.000889	0.106***	-	-	-	-	-	-	-	-	-	-	-	-	-		
tothhexp	-1.07e-05***	-6.15e-06***	-1.46e-05***	-4.33E-06	-2.52e-05*	-4.33E-05	1.35E-06	2.59E-07	-1.71E-07	5.04e-06**	0.00482	6.35E-05	0.000249 ***	5.04E-05	0.000105			
fhheduc	0.00145	0.00294	-0.00021	0.0780** *	0.0391***	0.170***	0.00485** *	0.00449* **	0.00646** *	0.00541**	-0.0155	0.00020 2	0.000184	-0.158**	0.000572			
mhheduc	-0.002	-0.0014	-0.00264*	0.0622** *	0.0373***	0.118**	-0.00197	-0.000177	-0.000605	-0.00178	0.0374***	0.00123 **	0.000161	0.0796	-0.000886			
fsalaried	-	-	-	0.0327** *	-0.0422***	-0.0273**	-0.346**	-0.218**	-0.779**	0.0190*	0.0202*	-0.00229	0.0320**	-0.417***	0.0137* **	-0.00102	0.0751	0.00382
fcontp	0.0183** *	-0.0245***	-0.0135***	0.271***	0.172***	0.662***	-0.00458	-0.00627	-0.0276***	0.000705	-0.154***	0.00967 ***	0.00244* **	-0.399**	0.0254** *			
fnowork	-	-	-	0.0239** *	-0.0312***	-0.0179**	-0.107	-0.0246	-0.222	0.0111	0.00713	-0.00302	0.0178*	-0.178**	0.0106* **	0.00129*	-0.119	0.0184** *
msalaried	0.00982	0.0244	-0.00544	0.228	0.237***	0.644**	-0.0305**	0.0462** *	-0.0232*	-	0.0492***	-0.0634	-0.0043	0.00166	-0.261	-0.000531		
mcontp	0.0236** *	0.0208*	0.0284***	0.375**	0.394***	0.728***	-	-	-	-	-	-	-	-	-	-		
mnwork	0.0281** *	0.0484***	0.00808	-0.185	-0.139	-0.238	-0.0258	-0.026	-0.0021	-0.0355*	-0.333***	-0.00474	-0.00016	-1.009**	-0.0168**			
adulteq	0.00134	0.000188	0.00297*	0.0546**	0.0102	0.191***	0.00415**	0.00653* **	0.00685** *	-0.00125	-0.0647***	-0.0521	-0.124***	-0.308***	-0.508**			
nkidslt5	-0.000758	-0.0023	0.000565	0.0526*	0.0326*	0.0504	0.00217	0.00204	-0.000838	0.00745**	0.0526**	-0.323**	0.0178	0.0578	-0.278			
nkids1220	-0.00293	-0.00389	-0.00283	0.0317	0.0299	0.1	-0.00521*	0.00717* *	-0.00680**	-0.0013	-0.00132	-0.154	0.0181	0.0777	-0.0814			
nummeals	0.00333	0.0217***	-0.0141***	0.586***	0.308***	1.316***	-	-	-	-	-	-	-	-	-			
Constant	4.001***	3.337***	3.249***	3.585***	2.385***	4.468***	0.440***	0.514***	0.435***	0.548***	2.343***	9.931** *	4.851***	9.612***	26.01***			
Observations	10,875	10,875	10,875	10,875	10,875	10,875	10,875	10,875	10,875	10,875	10,875	10,875	10,875	10,875	10,875			
R-squared	0.085	0.017	0.19	0.171	0.1	0.236	0.059	0.053	0.11	0.061	0.035	0.028	0.031	0.021	0.035			

*** p<0.01, ** p<0.05, * p<0.1; green highlighting is a significantly positive effect, orange highlighting is a significantly negative effect

Annex Table 6.12 Food Expenditure Share Regressions

VARIABLES	cerealsr	rootshr	nutshr	vegshr	frushr	fishshr	mlkshr	eggshr	bevshr	meatshr	sugshr	oilshr	miscshr
hgender	0.0409**	0.00167	0.00618	0.0162**	-0.0131	-0.0197**	-0.000143	0.000855	5.46E-05	0.0204***	-0.00387*	-0.00616**	-0.00254*
hage	-0.000204**	0.000318**	-1.62E-05	3.22E-05	-3.56E-05	2.19E-05	-6.03E-07	-6.50e-06***	-8.88e-06**	-4.05e-05*	-1.15e-05**	-3.64e-05***	-1.25e-05***
1.citysize	0.0198**	-0.0305***	0.00844**	-0.00369	0.0150***	0.0214***	5.30E-05	-0.000562	0.00412**	-0.00181	0.00628***	0.00727***	0.00106**
2.citysize	-0.0224*	0.0187**	-0.0135**	0.0353***	-0.00374	0.0540***	-0.00053	-0.000165	-0.00254	-0.00979	0.00482**	0.00955***	0.000874
3.citysize	-0.0876***	0.0690***	-0.0183**	0.0373***	0.0175***	0.00407	0.00254	0.00267	0.0267***	-0.0105	0.000324	0.00232	0.0285***
shareownfarm	-0.151***	0.158***	0.0433***	-0.110***	0.00333	0.0965***	0.000198	0.00171	0.00458	0.0413**	-0.000798	0.00979*	-0.0096
OwnfarmShare ownfarm	0.0953***	0.0680***	0.0297***	0.00432	-0.00351	0.0922***	-0.00128	-0.00103	0.0165***	0.0390***	0.00994***	-0.0263***	-0.00761**
totexp	0.00409***	0.00419**	0.00220**	0.00235**	0.000709*	0.00202**	0.000295*	0.000194**	0.00289**	0.00414**	0.000354**	0.000281**	0.000497**
fhheduc	-0.00545**	0.00198	-0.000297	-0.000452	-0.000162	0.000935	0.000202*	0.000211	-0.00063	0.00348**	0.000598*	0.000144	-0.000557*
mhheduc	-0.000205	-0.000795	0.00378**	-0.00182*	-0.00156	0.000535	7.98E-05	0.000401**	0.00296**	0.00294*	-4.90E-05	0.000938**	0.000345
fsalaried	-0.0359**	0.00592	-0.00644	0.0160*	0.00275	0.0168*	0.00546**	-0.0014	0.00154	0.00382	-0.00405*	-0.00727**	0.00282
fcontp	-0.0261***	0.00343	-0.00356	0.00632*	0.00286	0.0197***	3.68E-05	0.00220***	-0.000941	-0.00505	0.00318***	0.00264**	-0.000276
fnowork	-0.0221	0.00916	-0.00221	-0.000646	0.00233	0.0155**	0.00176	-0.000895	0.00848**	0.00673	0.000604	-0.00153	-0.000233
msalaried	0.0231	-0.0129	-0.00213	-0.00475	-0.0178	0.0164**	0.000788	-0.0004	0.0027	-0.00314	-0.000932	-0.000653	-0.000268
mcontp	0.0404**	0.00539	0.00168	-0.00763	-0.0256*	0.0106	6.16E-05	-0.000609	-0.00166	-0.011	-0.00237	0.00664***	-0.0027
mnowork	0.00117	-0.0247	0.0319**	0.0240*	-0.00449	-0.0226**	0.000537	0.00235	-0.00328	0.00477	0.00248	-0.00873*	-0.00336
nkidslt5	0.00917**	-0.00724**	0.00641**	-0.00112	0.00345	-0.0028	0.000309*	0.000578**	0.000577	0.00446**	0.000398	3.98E-05	-0.000265
nkids1220	0.00717**	-0.00513	0.00410**	0.000663	-0.000446	-0.000214	0.000256*	-0.000121	-0.000449	0.00284**	0.000512	-0.000123	0.000854**
nummeals	0.0206***	-0.0186***	-0.00456	0.0150***	-0.00171	0.00336	0.000176	0.00141***	0.000457	0.0108***	0.00308***	0.00237***	0.00242***
Constant	0.304***	0.0211	0.164***	0.236***	0.117***	0.0660***	-0.000668	-0.000491	0.00843	0.00948	0.0129***	0.0308***	0.0318***
Observations	10,873	10,873	10,873	10,873	10,873	10,873	10,873	10,873	10,873	10,873	10,873	10,873	10,873
R-squared	0.021	0.105	0.022	0.051	0.009	0.068	0.035	0.011	0.113	0.053	0.032	0.068	0.109

*** p<0.01, ** p<0.05, * p<0.1; green highlighting is a significantly positive effect, orange highlighting is a significantly negative effect

Annex Table 6.13 Monteiro et al's food groupings (2010)

Table 1

Food classification based on the extent and purpose of industrial processing.

Food group	Extent and purpose of processing	Examples *
Group 1: unprocessed or minimally processed foods	No processing, or mostly physical processes used to make single whole foods more durable, accessible, convenient, palatable, or safe	Fresh, chilled, frozen, vacuum-packed fruits, vegetables, fungi, roots and tubers; grains (cereals) in general; fresh, frozen and dried beans and other pulses (legumes); dried fruits and 100% unsweetened fruit juices; unsalted nuts and seeds; fresh, dried, chilled, frozen meats, poultry and fish; fresh and pasteurized milk, fermented milk such as plain yoghurt; eggs; teas, coffee, herb infusions, tap water, bottled spring water
Group 2: processed culinary or food industry ingredients	Extraction and purification of components of single whole foods, resulting in producing ingredients used in the preparation and cooking of dishes and meals made up from Group 1 foods in homes or traditional restaurants, or else in the formulation by manufacturers of Group 3 foods	Vegetable oils, margarine, butter, milk cream lard; sugar, sweeteners in general; salt; starches, flours, and "raw" pastas and noodles (made from flour with the addition only of water); and food industry ingredients usually not sold to consumers as such, including high fructose corn syrup, lactose, milk and soy proteins, gums, and preservatives and cosmetic additives
Group 3: ultra-processed food products	Processing of a mix of Group 2 ingredients and Group 1 foodstuffs in order to create durable, accessible, convenient, and palatable ready-to-eat or to-heat food products liable to be consumed as snacks or desserts or to replace home-prepared dishes	Breads, biscuits (cookies), cakes and pastries; ice cream; jams (preserves); fruits canned in syrup; chocolates, confectionery (candies), cereal bars, breakfast cereals with added sugar; chips, crisps; sauces; savoury and sweet snack products; cheeses; sugared fruit and milk drinks and sugared and "no-cal" cola, and other soft drinks; frozen pasta and pizza dishes; pre-prepared meat, poultry, fish, vegetable and other "recipe" dishes; processed meat including chicken nuggets, hot dogs, sausages, burgers, fish sticks; canned or dehydrated soups, stews and pot noodle, salted, pickled, smoked or cured meat and fish; vegetables bottled or canned in brine, fish canned in oil; infant formulas, follow-on milks, baby food

* These listings do not include alcoholic drinks. The examples given are not meant to be complete. Many others can be added, especially to group 3, using the general principles specified in the text and as indicated in the second column.

7. Bibliography

- Armar-Klimesu, M. (2000). Urban agriculture and food security, nutrition and health. *Growing Cities, Growing Food. Urban Agriculture on the Policy Agenda*, 99–118. Retrieved from <http://www.ruaf.org/sites/default/files/Theme4.PDF>
- Bellows, A. C., Brown, K., & Smit, J. (2003). Health Benefits of Urban Agriculture. *Community Food*. <https://doi.org/10.2105/AJPH.19.6.702-b>
- Bermudez, O. I., Lividini, K., Smitz, M. F., & Fiedler, J. L. (2012). Estimating micronutrient intakes from Household Consumption and Expenditures Surveys (HCES): an example from Bangladesh. *Food and Nutrition Bulletin*, 33(3 Suppl), 208–213.
- Berti, P. R. (2012). Intra-household distribution of food: a review of the literature and discussion of the implications for food fortification programs. *Food and Nutrition Bulletin*, 33(3 Suppl).
- Coates, J., Colaiezzi, B., Fiedler, J. L., Wirth, J., Lividini, K., & Rogers, B. (2012). A program needs-driven approach to selecting dietary assessment methods for decision-making in food fortification programs. *Food and Nutrition Bulletin*, 33(3 Suppl), 146–156.
- Darmon K, N. D. (2008). Does social class predict diet quality? *Am J Clin Nutr*.
- Dary, O., & Jariseta, Z. R. (2012). Validation of dietary applications of Household Consumption and Expenditures Surveys (HCES) against a 24-hour recall method in Uganda. *Food and Nutrition Bulletin*, 33(3 Suppl), 190–198.
- De Weerd, J., Beegle, K., Friedman, J., Gibson, J., & Bank, W. (2014). The Challenge of Measuring Hunger, (January), 1 online resource (28 p.). <https://doi.org/10.1596/1813-9450-6736>
- Delisle, H. (2010). Findings on dietary patterns in different groups of {African} origin undergoing nutrition transition. *Applied Physiology Nutrition and Metabolism-Physiologie Appliquee Nutrition Et Metabolisme*, 35(2), 224–228. <https://doi.org/10.1139/H10-008>
- Delisle, H. F., Vioque, J., & Gil, A. (2009). Dietary patterns and quality in West-African immigrants in Madrid. *Nutrition Journal*, 8, 3. <https://doi.org/10.1186/1475-2891-8-3>
- Dop, M. C., Pereira, C., Mistura, L., Martinez, C., & Cardoso, E. (2012). Using Household Consumption and Expenditures Survey (HCES) data to assess dietary intake in relation to the nutrition transition: a case study from Cape Verde. *Food and Nutrition Bulletin*, 33(3 Suppl), 221–227.
- Fiedler, J. L. (2013). Towards overcoming the food consumption information gap: Strengthening household consumption and expenditures surveys for food and nutrition policymaking. *Global Food Security*, 2(1), 56–63. <https://doi.org/10.1016/j.gfs.2012.09.002>
- Fiedler, J. L., Carletto, C., & Dupriez, O. (2012). Still waiting for Godot? Improving Household Consumption and Expenditures Surveys (HCES) to enable more evidence-based nutrition policies. *Food and Nutrition Bulletin*, 33(3 Suppl), 242–251.
- Fiedler, J. L., Lividini, K., Bermudez, O. I., & Smitz, M. F. (2012). Household Consumption and Expenditures Surveys (HCES): a primer for food and nutrition analysts in low- and middle-income countries. *Food and Nutrition Bulletin*, 33(3 Suppl), 170–184.

- Fiedler, J. L., Lividini, K., Zulu, R., Kabaghe, G., Tehinse, J., & Bermudez, O. I. (2013). Identifying Zambia's industrial fortification options: Toward overcoming the food and nutrition information gap-induced impasse. *Food and Nutrition Bulletin*, 34(4), 480–500. <https://doi.org/00000034/00000004/art00012>
- Foeken, D., & Mwangi, A. (2000). Increasing food security through urban farming in Nairobi. *Growing Cities, Growing Food: Urban Agriculture on the Policy Agenda. A Reader on Urban Agriculture*, 2(Kenya 1994), 303–327. Retrieved from <http://www.ruaf.org/sites/default/files/Nairobi.PDF>
- Frega, R., Lanfranco, J. G., De Greve, S., Bernardini, S., Geniez, P., Grede, N., ... de Pee, S. (2012). What linear programming contributes: world food programme experience with the "cost of the diet" tool. *Food and Nutrition Bulletin*, 33(3 Suppl), S228-34. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/23193775>
- Fung, T., McCullough, M., Newby, PK, Manson, J., Meigs, J., Rifai, N., Willett, W., Hu, F. (2005) Diet-quality scores and plasma concentrations of markers of inflammation and endothelial dysfunction. *American Journal of Clinical Nutrition*, 82: 163-73.
- Gaiha, R., Kaicker, N., Imai, K., Kulkarni, V., & Thapa, G. (2012). Has Dietary Transition Slowed Down in India: An analysis based on 50th, 61st and 66th Rounds of NSS. Rome: IFAD, 1–23. <https://doi.org/10.4337/9781781004296.00014>
- Gibson, R. S., & Cavalli-Sforza, T. (2012). Using reference nutrient density goals with food balance sheet data to identify likely micronutrient deficits for fortification planning in countries in the Western Pacific region. *Food and Nutrition Bulletin*, 33(3 Suppl), 214–220.
- Huijbregts, P., Feskens, E., Räsänen, L., Fidanza, F., Nissinen, A., Menotti, A., & Kromhout, D. (1997). Dietary pattern and 20 year mortality in elderly men in Finland, Italy, and The Netherlands: longitudinal cohort study. *Bmj*, 315(7099), 13-17.
- Imamura, F., Micha, R., Khatibzadeh, S., Fahimi, S., Shi, P., Powles, J., & Mozaffarian, D. (2015). Dietary quality among men and women in 187 countries in 1990 and 2010: A systematic assessment. *The Lancet Global Health*, 3(3), e132–e142. [https://doi.org/10.1016/S2214-109X\(14\)70381-X](https://doi.org/10.1016/S2214-109X(14)70381-X)
- Imhoff-Kunsch, B., Flores, R., Dary, O., & Martorell, R. (2012). Methods of using household consumption and expenditures survey (HCES) data to estimate the potential nutritional impact of fortified staple foods. *Food and Nutrition Bulletin*, 33(3 Suppl), 185–189.
- Instituto Nacional de Estatística (2010). IOF 2008/09 Summary Report from a series of missions to the National Statistical Institute of Mozambique, Maputo Mozambique.
- Jarisseta, Z. R., Dary, O., Fiedler, J. L., & Franklin, N. (2012). Comparison of estimates of the nutrient density of the diet of women and children in Uganda by Household Consumption and Expenditures Surveys (HCES) and 24-hour recall. *Food and Nutrition Bulletin*, 33(3 Suppl), 199–207.
- Kim, S., Haines, P. S., Siega-Riz, A. M., & Popkin, B. M. (2003). The Diet Quality Index-International (DQI-I) provides an effective tool for cross-national comparison of diet quality as illustrated by China and the United States. *The Journal of Nutrition*, 133(11), 3476–3484.

- Korkalo, L., Hauta-alus, H., & Mutanen, M. (2011). Food composition tables for Mozambique. *Helsinki, Finland: University of Helsinki*.
- Lim, S. S., Vos, T., Flaxman, A. D., Danaei, G., Shibuya, K., Adair-Rohani, H., Ezzati, M. (2012). A comparative risk assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters in 21 regions, 1990-2010: A systematic analysis for the Global Burden of Disease Study 2010. *The Lancet*, 380(9859), 2224–2260. [https://doi.org/10.1016/S0140-6736\(12\)61766-8](https://doi.org/10.1016/S0140-6736(12)61766-8)
- Mabli, J., Castner, L., Ohls, J., Fox, M.K., Crepinsek, M.K., Condon, E. (2010). Food Expenditures and Diet Quality Among Low-Income Households and Individuals.
- Madaleno, I. (2000). Urban agriculture in Belem, Brazil. *Cities*, 17(1), 73–77. [https://doi.org/10.1016/S0264-2751\(99\)00053-0](https://doi.org/10.1016/S0264-2751(99)00053-0)
- Maxwell, D. G. (1995). Alternative Food Security Strategy : A Household Analysis of Urban Agriculture in Kampala, 23(10), 1669–1681.
- McCullough, M. L., Feskanich, D., Stampfer, M. J., Giovannucci, E. L., Rimm, E. B., Hu, F. B., ... Willett, W. C. (2002). Diet quality and major chronic disease risk in men and women: Moving toward improved dietary guidance. *American Journal of Clinical Nutrition*, 76(6), 1261–1271.
- McCullough, M. L., & Willett, W. C. (2006). Evaluating adherence to recommended diets in adults: the Alternate Healthy Eating Index. *Public Health Nutrition*, 9(1A), 152–157. <https://doi.org/10.1079/PHN2005938>
- Micha, R., Kalantarian, S., Wirojratana, P., Byers, T., Danaei, G., Elmadfa, I., ... Mozaffarian, D. (2012). Estimating the global and regional burden of suboptimal nutrition on chronic disease: methods and inputs to the analysis. *European Journal of Clinical Nutrition*, 66(1), 119–129. <https://doi.org/10.1038/ejcn.2011.147>
- Micha, R., Khatibzadeh, S., Shi, P., Fahimi, S., Lim, S., Andrews, K. G., ... Mozaffarian, D. (2014). Global, regional, and national consumption levels of dietary fats and oils in 1990 and 2010: a systematic analysis including 266 country-specific nutrition surveys. *BMJ (Clinical Research Ed.)*, 348(apr14_18), g2272. <https://doi.org/10.1136/bmj.g2272>
- Monteiro, C., Levy, R., Claro, R., De Castro, I., & Cannon, G. (2010). A new classification of foods based on the extent and purpose of their processing. *Cadernos de Saude Publica / Ministerio Da Saude, Fundacao Oswaldo Cruz, Escola Nacional de Saude Publica*, 26(11), 2039–2049. <https://doi.org/10.1590/S0102-311X2010001100005>
- Murphy, S., Ruel, M., & Carriquiry, A. (2012). Should Household Consumption and Expenditures Surveys (HCES) be used for nutritional assessment and planning? *Food and Nutrition Bulletin*, 33(3 Suppl), 235–241.
- National Directorate for Studies and Policy Analysis (DNEAP) (Mozambique), National Institute of Statistics (INE) (Mozambique), United Nations Population Fund (UNFPA). Mozambique Household Budget Survey 2008-2009.
- Nelson, C. (2010). Don't mourn, organize. *Academy*, 96(1), 10–14. <https://doi.org/10.1002/jid>

- Neufeld, L. M., Dary, O., Macdonald, B., & Harding, K. B. (2012). Measurement of Food Consumption to Inform Food Fortification and other Nutrition Programs: Methods and Applications. *Food and Nutrition Bulletin*, 33(3 (supplement)), 141–145.
- Popkin, B. M. (2014). Nutrition, agriculture and the global food system in low and middle income countries. *Food Policy*, 47, 91–96. <https://doi.org/10.1016/j.foodpol.2014.05.001>
- Popkin, B. M., & Gordon-Larsen, P. (2004). The nutrition transition: worldwide obesity dynamics and their determinants. *International Journal of Obesity and Related Metabolic Disorders : Journal of the International Association for the Study of Obesity*, 28 Suppl 3(Suppl 3), S2-9. <https://doi.org/10.1038/sj.ijo.0802804>
- Powles, J., Fahimi, S., Micha, R., Khatibzadeh, S., Shi, P., Ezzati, M., ... Mozaffarian, D. (2013). Global, regional and national sodium intakes in 1990 and 2010: a systematic analysis of 24 h urinary sodium excretion and dietary surveys worldwide. *BMJ Open*, 3(12), e003733. <https://doi.org/10.1136/bmjopen-2013-003733>
- Rashid, D. A., Smith, L. C., & Rahman, T. (2011). Determinants of dietary quality: Evidence from Bangladesh. *World Development*, 39(12), 2221–2231. <https://doi.org/10.1016/j.worlddev.2011.05.022>
- Reardon, T., Tschirley, D., Minten, B., Haggblade, S., Liverpool-Tasie, S., & Dolislager, M. (2015). Transformation of African Agrifood Systems in the New Era of Rapid Urbanization and the Emergence of a Middle Class. Chapter in the Proceedings Volume of the ReSAKSS Annual Conference, “Beyond a Middle Income Africa,” Trends and Outlook Report Conference Held in Addis Ababa, September 1-3, 2015., 1–16.
- Ruel, M. T. (2003). Is dietary diversity an indicator of food security or dietary quality? A review of measurement issues and research needs. *Food and Nutrition Bulletin*, 24(2), 231–232.
- Smith, L. C., & Subandoro, A. (2007). Measuring Food Security Using HH Expenditure Surveys. <https://doi.org/10.2499/0896297675>
- Sodjinou, R., Agueh, V., Fayomi, B., & Delisle, H. (2009). Dietary patterns of urban adults in Benin: relationship with overall diet quality and socio-demographic characteristics. *European Journal of Clinical Nutrition*, 63, 222–228. <https://doi.org/10.1038/sj.ejcn.1602906>
- Tschirley, D. L., Snyder, J., Dolislager, M., Reardon, T., Haggblade, S., Goeb, J., Meyer, F. (2015). Africa’s unfolding diet transformation: implications for agrifood system employment. *Journal of Agribusiness in Developing and Emerging Economies*, 5(2), 0–48. <https://doi.org/10.1108/JADEE-01-2015-0003>
- Tschirley, D., Reardon, T., Dolislager, M., Snyder, J. (2015). The Rise of a Middle Class in East and Southern Africa: Implications for Food System Transformation. *Journal of International Development*, 27, 628-646.
- US Department of Agriculture, Agricultural Research Service, Nutrient Data Laboratory. USDA National Nutrient Database for Standard Reference, Release 28. Version Current: September 2015. Internet: [/nea/bhnrc/ndl](http://nea/bhnrc/ndl)

- Vecchia, C. La, & Majem, L. S. (2015). Evaluating trends in global dietary patterns. *The Lancet Global Health*, 3(3), e114–e115. [https://doi.org/10.1016/S2214-109X\(15\)70011-2](https://doi.org/10.1016/S2214-109X(15)70011-2)
- Vorster, H. H., Kruger, A., & Margetts, B. M. (2011). The nutrition transition in Africa: can it be steered into a more positive direction? *Nutrients*, 3(4), 429–441. <https://doi.org/10.3390/nu3040429>
- Weisell, R., & Dop, M. C. (2012). The adult male equivalent concept and its application to Household Consumption and Expenditures Surveys (HCES). *Food and Nutrition Bulletin*, 33(3 Suppl), 157–162. <https://doi.org/10.1177/15648265120333S203>
- World Health Organization, W. (2015). Population nutrient intake goals for preventing diet-related chronic diseases. *Www.Who.Int*, 346(12), 871–876.
- World Health Organization. (2003). Diet, nutrition, and the prevention of chronic diseases: report of a joint WHO/FAO expert consultation (Vol. 916). World Health Organization.
- Zeza, A., & Tasciotti, L. (2010). Urban agriculture, poverty, and food security: Empirical evidence from a sample of developing countries. *Food Policy*, 35(4), 265–273. <https://doi.org/10.1016/j.foodpol.2010.04.007>
- Ziraba, A. K., Fotso, J. C., & Ochako, R. (2009). Overweight and obesity in urban Africa: A problem of the rich or the poor? *BMC Public Health*, 9, 465–473. <https://doi.org/10.1186/1471-2458-9-465>