

# The Vanishing U. S. Cattle Cycle: A Stochastic Cycle Approach

Yunhan Li and J. Scott Shonkwiler

We re-examine the existence of cattle cycles based on U.S. beef cow inventories from 1979 to 2019. Our analysis begins with a basic first-order stochastic cycle model and finds a cattle cycle of 16.54 years, significantly longer than the presumed 10- to 12-year cycle. Typically cycles become longer before they disappear. Upon further investigation, we re-estimate the length of the cycle by applying a second-order stochastic cycle model which improves goodness-of-fit. Surprisingly, the cycle length is estimated to be infinitely long, implying the complete disappearance of the beef cow cycle.

*Key words:* Cattle Cycle, Cattle Dynamics, Cattle Inventory, Stochastic Cycle

## Introduction

The dynamics of beef cattle supply and the existence of cattle cycles have been widely researched topics in the last four decades. Jarvis (1974) was first to treat beef cows in the context of capital goods and recognized that increasing beef prices can actually lead to reduced slaughter in the short run. This study influenced empirical approaches to modeling the beef cattle herd such as formulated by Rucker, Burt, and LaFrance (1984) and stimulated a theoretical treatment of the dynamics of livestock production by Rosen (1987). Rosen, Murphy, and Scheinkman (1994) specifically addressed the existence of cattle cycles. More recently Aadland (2004) constructed a model to describe the putative 10-year cattle cycle by assuming that producers maximize a discounted stream of future profits subject to biological constraints and market forces.

Under a fully rational expectations theory, forward-looking ranchers should anticipate future cattle cycles, which were known to be approximately 10 years, and take actions to mitigate them. Of course, under the animal spirits paradigm (where emotions affect economic decisions) their actions could endogenously generate or realize another 10-year cycle in the future. There are also quite a few studies that put in question the appropriateness of fully rational expectations in the cattle industry (Nerlove and Fornari, 1998; Baak 1999; Chavas 2000) as most cattle market participants are found to be not fully rational (e.g. quasi-rational or boundedly rational). However, Aadland (2004) showed that individual optimizing behaviour endogenously generated approximate 10-year cycles in cattle supply even under a less restrictive assumption that ranchers are only boundedly rational (i.e. ranchers have limited access to future information and form expectations of future prices based on only past or current information). As will be seen in Section 2, the extraordinary change in the beef cow cycle suggests that previous studies of the cattle cycle may need to be re-evaluated.

Since it is difficult to visualize a persistent cycle from the recent semi-annual or annual data, we turn to the business cycle literature because periodic patterns in business cycles are typically not easily discerned. In 1927, Eugen Slutsky published a paper (subsequently available in English

---

Yunhan Li (corresponding author, [jss1@uga.edu](mailto:jss1@uga.edu)) and J. Scott Shonkwiler are former doctoral student and adjunct professor in the Department of Agricultural and Applied Economics at The University of Georgia.

This work was supported in part by Hatch Project GEO00793 [Project Accession no. 1009593] from the USDA National Institute of Food and Agriculture. The comments of Jeff Dorfman, Wenying Li, Chen Zhen, the editor, and reviewers are gratefully acknowledged.

This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License.

Review coordinated by Darren Hudson.

in 1937) which was the first to introduce the notion of a stochastic cycle. He posited that under certain circumstances the summation of random causes may be the source of a cyclical process. He also noted that cycles may transition over time so as to make it difficult to detect a regular pattern. Slutsky (1927) was able to closely approximate a 19<sup>th</sup> Century English business cycle using his stochastic model. Beveridge and Nelson (1981) gave a review of business cycle analyses and their study was one of the first to recognize and account for the non-stationarity of macroeconomic data. Harvey (1985) developed a stochastic cycle model which allowed the amplitude and phase of the cycle to vary over time. Harvey divided his datasets of macroeconomic series into two periods. He found that before 1948, a six to nine year cycle was evident in GNP. With more recent data, Harvey was less confident that cyclical components could be identified. A highly influential paper by Howitt and McAfee (1992) addressed the idea that the business cycle results from exogenous shocks that lead to economic fluctuations in an otherwise stable system. They constructed a rational expectations-animal spirits cycle. The notion of animal spirits relates to random waves of optimism or pessimism that are not necessarily based on economic conditions. Howitt and McAfee proposed that "When spirits are high, firms expect a high level of employment, and hence a high level of aggregate demand... This encourages firms to hire more aggressively, thus validating the original expectation." (p. 494). They showed that the animal spirits cycle may be stable under Bayesian learning: "If people start with diffuse priors, there is a positive probability that accidental correlations during the early stages of learning could lead them forever into the self-fulfilling beliefs of the animal spirits cycle." (p. 505). Nevertheless, they did conclude that the probability of a perpetual cycle seemed remote. In 1997, Hodrick and Prescott introduced a low-pass filter to de-trend economic time series. If properly de-trended, the cycle then is the residual of the series. The use of model based time series filters to extract cyclical components was reconsidered by Harvey and Trimbur (2003). They extended the original trend plus cycle model of Harvey (1985) to allow for higher ordered cycles. They showed that a second order stochastic cycle model tends to fit the data better and can provide a smoother (i.e. less noisy) cyclical representation. Based on their experience the second order model should also yield more plausible values for the cycle length. To the best of our knowledge, this is the first study that applies business cycle models to depict the cyclical pattern in the U.S. beef cattle inventory.

This article is outlined as follows: In Section 2, we describe the data applied in this study. We use the first- and second-order stochastic models to fit U.S. beef cow inventory data in Sections 3 and 4, respectively, where estimation results and their implications are also provided. In the last Section 5, we draw conclusions and provide further discussion.

## Data Description

The data used in this study is sourced from National Agricultural Statistics Service<sup>1</sup> (NASS) of the U.S. Department of Agriculture (USDA), which reports the U.S. beef cow inventory twice a year – numbers are reported in every January and July. While the beef cow inventory of January is based on survey data, the mid-year numbers are derived from marketing information and a smaller survey. Figure 1 plots annual beef cow inventory numbers corrected for mean and trend<sup>2</sup> for the 62-year period from 1920 to 1981. It is evident that generally numbers peaked mid-decade and tended to bottom out around the beginning of each decade—suggesting a ten year cattle cycle. In fact, when a trigonometric regression with  $\sin(\lambda t)$  and  $\cos(\lambda t)$  along with a polynomial in time is performed over this period,  $\lambda$  is estimated to be 0.6235, which implies a cycle length of 10.08 years.

---

<sup>1</sup> <https://quickstats.nass.usda.gov/>

<sup>2</sup> A linear trend has been removed from the original series.

However, Figure 1 stands in contrast to the trend and seasonally adjusted<sup>3</sup> semi-annual beef cow inventories for the recent 40 years (1979-2019) depicted in Figure 2. Since 1995, the number of beef cows (trend and seasonally adjusted) appears to have little cyclicity. We do need to mention that there are two missing values for this series occurring mid-year of 2013 and 2016. These values were calculated by estimating an autoregressive model of order four with constant, seasonal dummy, and trend and performing a grid search over the missing values such that the model's sum of squared errors was minimized. The resulting estimates of beef cow numbers were 29.9 million for 2013 and 31.4 million for 2016. The little cyclical pattern observed in the more recent beef cow cycle—whether it is significantly longer than ten years or has disappeared—calls for the need to re-examine the cattle cycle.

### The Basic Stochastic Cycle Model

This study begins with analysing the semi-annual beef cow inventory data using the Harvey and Trimbur (2003) approach, which is referred to as the Basic Stochastic Cycle Model in this study, to attempt to estimate the length of the beef cow cycle. The irregular patterns evidenced in Figure 2 suggest that the amplitudes and phases of this series have been evolving over time. To address these issues we investigate a random walk plus cycle model which allows i) analysis of non-stationary data; ii) direct estimation of cycle length; and iii) shifting phase and changing amplitudes.

The model is specified to be a random walk (with drift) with a stochastic cycle (Harvey, 1989) for the stochastic series  $y_t$ .

$$\begin{aligned} (1) \quad & y_t = \mu_t + \psi_t + x_t \delta + \varepsilon_t \\ (2) \quad & \mu_t = \mu_{t-1} + \beta + \eta_t \\ (3) \quad & \psi_t = \rho \{ \cos(\lambda) \psi_{t-1} + \sin(\lambda) \psi_{t-1}^* \} + \kappa_t \\ (4) \quad & \psi_t^* = \rho \{ -\sin(\lambda) \psi_{t-1} + \cos(\lambda) \psi_{t-1}^* \} + \nu_t \end{aligned}$$

Here  $\mu_t$  and  $\psi_t$  represent dynamic unobservables associated with a random walk (with drift  $\beta$ ) and a cyclical process (with frequency  $\lambda$ ), respectively.  $x_t$  is coded as one if the inventory number is reported in January. Otherwise  $x_t = 0$ . The coefficient  $\rho$  is termed the damping factor and  $\rho = 1$  if the cyclical process is non-stationary. The error processes  $\varepsilon_t$ ,  $\eta_t$ ,  $\kappa_t$ , and  $\nu_t$  are assumed to be iid normal with variance-covariance matrix

$$\begin{bmatrix} \sigma_\varepsilon^2 & 0 & 0 & 0 \\ 0 & \sigma_\eta^2 & 0 & 0 \\ 0 & 0 & \sigma_\kappa^2 & 0 \\ 0 & 0 & 0 & \sigma_\nu^2 \end{bmatrix}$$

The initial conditions  $\psi_0$  and  $\psi_0^*$  determine the initial amplitude and phase shift of the series and  $\mu_0$  denotes the initial level of the series which can be estimated in lieu of fitting a constant to the model. Parker and Shonkwiler (2014) showed that the reduced form of the model can be written in terms of the observable process  $y_t$ , the unknown parameters, and the error processes:

$$\begin{aligned} (1) \quad \Delta y_t = & 2\rho \cos \lambda \Delta y_{t-1} - \rho^2 \Delta y_{t-2} + \beta^* + \eta_t - 2\rho \cos \lambda \eta_{t-1} + \rho^2 \eta_{t-2} \\ & + \Delta \kappa_t - \rho \cos \lambda \Delta \kappa_{t-1} + \rho \sin \lambda \Delta \nu_{t-1} + \Delta \xi_t - 2\rho \cos \lambda \Delta \xi_{t-1} + \rho^2 \Delta \xi_{t-2} \end{aligned}$$

<sup>3</sup> A quadratic trend and a semi-annual dummy variable are included in the regression.

where  $\Delta$  denotes the (first) difference operator,  $\beta^*$  represents the sum of deterministic terms, and  $\xi_t = x_t\delta + \varepsilon_t$ . In time series parlance the series is a type of ARIMA(2,1,2) process. This representation is valid when  $y_t$  has a random walk component, i.e.  $\sigma_\eta^2 > 0$  and consequently the series must be first differenced to achieve stationarity. If this is not the case, the model with  $\sigma_\eta^2 = 0$  simplifies to

$$(6) \quad y_t = 2\rho\cos\lambda y_{t-1} - \rho^2 y_{t-2} + \beta t + \kappa_t - \rho\cos\lambda\kappa_{t-1} + \rho\sin\lambda v_{t-1} + \varepsilon_t \\ - 2\rho\cos\lambda\xi_{t-1} + \rho^2\xi_{t-2}$$

or a type of ARMA(2,2) process with constant trend. An additional simplification of the dynamic process occurs when all the noise in the system is due to the stochastic cycle. In this case  $\sigma_\varepsilon^2 = 0$ , and then

$$(7) \quad y_t = 2\rho\cos\lambda y_{t-1} - \rho^2 y_{t-2} + \beta t + \kappa_t - \rho\cos\lambda\kappa_{t-1} + \rho\sin\lambda v_{t-1} \\ - 2\rho\cos\lambda x_{(t-1)}\delta + \rho^2 x_{(t-2)}\delta.$$

### Estimation Results Model I

A MATLAB program using double sided numerical first and second derivatives of the likelihood function implied by the state-space model was used to obtain parameter estimates (Harvey, 1989 p. 126). Estimation results for the basic stochastic cycle model are reported in Table 1, where  $\mu_0$ ,  $\Psi_0$ , and  $\Psi_0^*$  represent the initial values for  $\mu_t$ ,  $\Psi_t$ , and  $\Psi_t^*$ , respectively.  $\delta$  is the coefficient on the semi-annual dummy variable. The other parameters are defined the same as in Equations (1) – (4) with dependent variable being semi-annual beef cow inventory from 1979 to 2019. Imposing the customary restriction (Harvey, p.39) that  $\sigma_\kappa^2 = \sigma_v^2$  for the cow herd size, we found that  $\sigma_\varepsilon^2$  converged to zero under maximum likelihood estimation of the stochastic cycle model. We adopted a logarithmic time trend<sup>4</sup> as it resulted in a marginally better fit than a linear counterpart; its coefficient ( $\beta$ ) is estimated to be positive and significant at the 5% level. The highly significant coefficient on the semi-annual dummy variable ( $\delta$ ) shows that the cow herd tends to be smaller by almost 700 thousand (herd size is expressed in units of 100,000) on January 1 than on July 1. The estimate of  $\lambda$  implies a cattle cycle of  $\frac{2\pi}{\lambda}$ , about 33.1 periods or 16.57 years. Using the delta method, we find an approximate asymptotic 95% confidence interval for the cycle length of 12.68 to 20.46 years. Clearly this is significantly longer than the presumed 10 year cycle. Further the estimated amplitude  $\hat{\rho} = 0.94$  suggests that the cycle does not decay much from period to period.

The statistically significant estimate of  $\sigma_\eta$  indicates that there is a non-stationary component to beef cow numbers. Therefore the reduced form of the model follows Equation 5, with  $\xi_t = x_t\delta$  and  $\delta = [\beta, \beta_{DV}]'$ . Here we also impose the restriction  $\psi_0 = \psi_0^*$ ; whereas it is often customary to set  $\psi_0 = \psi_0^* = 0$ . Clearly the initial condition for the cyclical component is significantly different than zero.

Robust standard errors (White, 1982) are calculated to ensure that inference is robust to distributional misspecification such as non-normality or heteroscedasticity; however the standard errors may be affected by autocorrelated errors. Therefore the Ljung-Box (1978) Q test (LBQ) for serially correlated residuals is calculated based on  $m=20$  lagged residual correlations and adjusted for degrees of freedom determined by  $m$  minus the number of parameters estimated excluding the constant,  $\mu_0$ , and the variance terms. The p-value of the test under the hypothesis

<sup>4</sup> The logarithmic time trend is sensitive to the value of the beginning time. We use  $t = 1$  to date the first observation that was observed in January 1979.

of temporally independent errors is shown to be 0.030 for Model I. Inspection of the correlogram then led to the inclusion of the stationary variable  $\Delta y_{t-3}$ .

*Estimation Results Model II*

The first order stochastic cycle model augmented with  $\Delta y_{t-3}$  has estimated parameters that are largely unchanged except for the coefficient on the logarithmic trend which becomes statistically insignificant. Interestingly,  $\lambda$  is estimated with more precision and the model now suggests a beef cow cycle of 16.54 years with a 95% confidence interval of 13.35 to 19.74 years. The variance of the cycle is  $\sigma_\psi^2 = \sigma_\kappa^2 / (1 - \rho^2) = 27.46$  which can be compared to the variance of the random walk which is  $t\sigma_\eta^2$  or 438.46 at the 80<sup>th</sup> observation. We cannot reject the hypothesis of independent errors at conventional levels based on the LBQ test calculated as above. In addition, the estimated amplitude ( $\hat{\rho} = 0.95$ ) does not vary much compared with that in Model I.

Were we to stop the analysis at this point, we would conclude that the augmented random walk model with stochastic trend demonstrates a substantial lengthening of the beef cattle cycle. The 95% interval for the cycle length does not include the ten to twelve year cycle that has been historically accepted. For this reason, we extend the analysis to a second order stochastic cycle representation to determine whether it offers an improvement in fit and corroborating evidence of a lengthening cattle cycle.

**The Higher-Order Stochastic Cycle Model**

*Generalized Stochastic Cycles*

Instead of the basic model in previous sections, which incorporates a first – order stochastic cycle, we apply a higher order stochastic model to explore the periodic dynamics of the beef cow inventories. Periodic behavior can be better identified by generalized cycles as they have more power concentrated near the spectral peak (Trimbur, 2006). Examples in Harvey and Trimbur (2003) showed smoother cycles in economic series can be extracted by generalized cyclical processes. Allowing for variation in the amplitude and phase over time as the first-order stochastic cycle does, the higher order cycles also allow the disturbances themselves to be periodic, thus producing a more flexible description of periodic behavior in time series data. Additionally, the interpretation of parameters remains straightforward and as such the cyclical pattern can be studied directly. Extensions to multivariable models are also possible.

The unobserved component  $\psi_t^{(m)}$  is defined as an  $m^{th}$ -order stochastic cycle, if

$$(8) \quad \begin{bmatrix} \psi_t^{(1)} \\ \psi_t^{(1)*} \end{bmatrix} = \rho \begin{bmatrix} \cos(\lambda) & \sin(\lambda) \\ -\sin(\lambda) & \cos(\lambda) \end{bmatrix} \begin{bmatrix} \psi_{t-1}^{(1)} \\ \psi_{t-1}^{(1)*} \end{bmatrix} + \begin{bmatrix} \kappa_t \\ \kappa_t^* \end{bmatrix}, t = 1, 2, \dots T$$

$$(9) \quad \begin{bmatrix} \psi_t^{(i)} \\ \psi_t^{(i)*} \end{bmatrix} = \rho \begin{bmatrix} \cos(\lambda) & \sin(\lambda) \\ -\sin(\lambda) & \cos(\lambda) \end{bmatrix} \begin{bmatrix} \psi_{t-1}^{(i)} \\ \psi_{t-1}^{(i)*} \end{bmatrix} + \begin{bmatrix} \psi_{t-1}^{(i-1)} \\ \psi_{t-1}^{(i-1)*} \end{bmatrix}, t = 1, 2, \dots T, i = 2, \dots m$$

where  $\kappa_t$  and  $\kappa_t^*$  are uncorrelated white-noise processes with zero mean and common variance  $\sigma_\kappa^2$ .  $\rho$  is the damping factor, which is bounded between 0 and 1. The cycle is stationary if  $\rho < 1$ . The stochastic nature of cycles comes from the disturbances  $\begin{bmatrix} \kappa_t \\ \kappa_t^* \end{bmatrix}$ . Higher order stochastic cycles are more flexible than the first order one because when order increases, the disturbances can impact cycles in different ways, thus resulting in different dynamic properties. For instance, a second-order stochastic cycle,  $\psi_t^{(2)}$ , has a first-order stochastic cycle  $\psi_{t-1}^{(1)}$  as a driving component, which allows the shocks to the second-order cycle to be periodic, as well.

Although cycles can be reduced to autoregressive moving-average (ARMA) form, they are more often represented in a state-space form, as this is the easiest way in terms of estimation and offers more intuitive parameterization than the reduced form. Therefore, we put an  $m^{\text{th}}$ -order cycle in a state-space form, with the transition equation specified as

$$(10) \quad \boldsymbol{\psi}_t = \mathbf{T}_m \boldsymbol{\psi}_{t-1} + \mathbf{c}_m \otimes \begin{bmatrix} \kappa_t \\ \kappa_t^* \end{bmatrix}$$

where  $\boldsymbol{\psi}_t = [\psi_t^{(m)}, \psi_t^{(m)*}, \psi_t^{(m-1)}, \psi_t^{(m-1)*}, \dots, \psi_t^{(1)}, \psi_t^{(1)*}]'$  is the state vector with  $2m$  elements. The last element of the  $m \times 1$  vector  $\mathbf{c}_m$  is one, with zeros elsewhere.  $\mathbf{T}_m$  is given by

$$(11) \quad \mathbf{T}_m = \mathbf{I}_m \otimes \mathbf{T} + \mathbf{S}_m \otimes \mathbf{I}_2$$

where the transition matrix  $\mathbf{T}$  is given by  $\mathbf{T} = \rho \begin{bmatrix} \cos(\lambda) & \sin(\lambda) \\ -\sin(\lambda) & \cos(\lambda) \end{bmatrix}$ , and  $\mathbf{S}_m$  is a matrix with ones on the diagonal strip to the right of the main diagonal and zeros elsewhere. The variance – covariance matrix of the disturbance vector  $\mathbf{c}_m \otimes \begin{bmatrix} \kappa_t \\ \kappa_t^* \end{bmatrix}$  is given by

$$(12) \quad \text{VAR} \left( \mathbf{c}_m \otimes \begin{bmatrix} \kappa_t \\ \kappa_t^* \end{bmatrix} \right) = \mathbf{c}_m \mathbf{c}_m' \otimes \begin{bmatrix} \sigma_\kappa^2 & 0 \\ 0 & \sigma_\kappa^2 \end{bmatrix}$$

where  $\text{VAR}(\cdot)$  forms the variance-covariance matrix.

#### *Fitting Annual Beef Cow Inventories using Second-Order Stochastic Cycle Model*

We then start analyzing this series with the random walk second—order stochastic cycle model. The system is specified in equations (13) – (16):

$$(13) \quad y_t = \mu_t + \psi_t^{(2)} + \log(t_t)\beta + x_t\delta + \varepsilon_t,$$

$$(14) \quad \mu_t = \mu_{t-1} + \eta_t,$$

$$(15) \quad \begin{bmatrix} \psi_t^{(1)} \\ \psi_t^{(1)*} \end{bmatrix} = \rho \begin{bmatrix} \cos(\lambda) & \sin(\lambda) \\ -\sin(\lambda) & \cos(\lambda) \end{bmatrix} \begin{bmatrix} \psi_{t-1}^{(1)} \\ \psi_{t-1}^{(1)*} \end{bmatrix} + \begin{bmatrix} \kappa_t \\ \kappa_t^* \end{bmatrix}, \text{ and}$$

$$(16) \quad \begin{bmatrix} \psi_t^{(2)} \\ \psi_t^{(2)*} \end{bmatrix} = \rho \begin{bmatrix} \cos(\lambda) & \sin(\lambda) \\ -\sin(\lambda) & \cos(\lambda) \end{bmatrix} \begin{bmatrix} \psi_{t-1}^{(2)} \\ \psi_{t-1}^{(2)*} \end{bmatrix} + \begin{bmatrix} \psi_{t-1}^{(1)} \\ \psi_{t-1}^{(1)*} \end{bmatrix}$$

where  $\mu_t$  is the trend component, and  $\psi_t^{(2)}$  is the  $2^{\text{nd}}$  – order stochastic cycle component. All parameters and disturbances follow the same definition as in equations (1) – (4), with the semi-annual dummy represented by  $x_t$ . We can also write equations (13) – (16) in the state-space form, with the measurement equation and transition equation specified as below:

$$(17) \quad y_t = \mathbf{Z}\boldsymbol{\alpha}_t + \mathbf{X}_t\boldsymbol{\beta} + \varepsilon_t \text{ (Measurement equation)}$$

$$(18) \quad \boldsymbol{\alpha}_t = \begin{bmatrix} \mu_t \\ \psi_t \end{bmatrix} = \begin{bmatrix} \mu_{t-1} + \eta_t \\ \mathbf{T}_2 \boldsymbol{\psi}_{t-1} + \mathbf{c}_2 \otimes \begin{bmatrix} \kappa_t \\ \kappa_t^* \end{bmatrix} \end{bmatrix} \text{ (Transition equation)}$$

where  $\mathbf{Z} = [1 \ 1 \ 0 \ 0 \ 0]$ , and  $\varepsilon_t$  is assumed to be iid normally distributed with mean zero and constant variance  $\sigma_\varepsilon^2$ .

The transition equation for the second order cyclical component is

$$T_2 = \begin{bmatrix} \rho \cos(\lambda) & \rho \sin(\lambda) & 1 & 0 \\ -\rho \sin(\lambda) & \rho \cos(\lambda) & 0 & 1 \\ 0 & 0 & \rho \cos(\lambda) & \rho \sin(\lambda) \\ 0 & 0 & -\rho \sin(\lambda) & \rho \cos(\lambda) \end{bmatrix}, c_2 = [0, 1]'$$

The system involves the nine parameters:  $\mu_0$ ,  $\beta$ ,  $\delta$ ,  $\rho$ ,  $\lambda$ ,  $\sigma_\varepsilon$ ,  $\sigma_\kappa = \sigma_{\kappa^*}$ ,  $\sigma_\eta$ , and the initial condition  $\psi_0 = \psi_0^*$  as with the first order cycle model. The system is estimated by maximum likelihood, and optimization is achieved by Kalman filter algorithm.

As mentioned in Section 2, given the concern that the results from using semi-annual data may partially be an artifact of the different survey used to calculate mid-year inventories, we use the second-order cycle model to fit annual beef cow inventory data, which includes the same 40 observations used earlier. The second order cycle estimation results are reported in Table 2. All the parameters have the same definitions as in Table 1. As the model iterated to convergence the estimate of  $\lambda$  became quite small. Because  $\lambda$  must be in the interval between zero and  $\pi$ , inclusive, it was replaced by its square to maintain non-negativity. Surprisingly the estimate of  $\lambda^2$  converged to zero—thus signifying an infinitely long cattle cycle. As with the first order cycle model the estimate of  $\sigma_\varepsilon$  also converged to zero. Because the LBQ statistic for the base model suggests some temporal non-independence of the error terms, we will focus on the model augmented with  $\Delta y_{t-3}$ .

Comparing the second order cycle augmented model to the first order cycle augmented model, we see that the logarithmic time trend has a coefficient that is now negative and statistically significant. The estimate of the damping factor,  $\rho$ , shrinks and the amount of the relative variation in the random components switches: the variance of the cycle is  $\sigma_\psi^2 = (1 + \rho^2)\sigma_\kappa^2 / (1 - \rho^2)^3 = 144.88$ , much greater than the variance of the first order cycle; while the variance of the random walk at the 80<sup>th</sup> observation is 235.16, much less than that of the first order cycle model. In choosing between models, we observe that the log likelihood and implied  $R^2$  of the second order model are both greater than those of the first order model, and this is accomplished with one less parameter. Further the very small value of the LBQ statistic (and consequently large p-value) indicates a much better representation of the data by the second order augmented model.

### Implications of $\lambda=0$

The fact that  $\lambda$  becomes zero in the second order cycle model implies the complete disappearance of the beef cow cycle. It may seem contradictory that a stochastic cycle model converges to a solution that indicates the absence of a cycle. To better understand the meaning of  $\lambda$  being equal to zero, consider the autocorrelation functions,  $\rho(\tau)$ , determined by the two stochastic cycle models. For the first order model  $\rho(\tau) = \rho^\tau \cos(\lambda\tau)$  and for the second order model the autocorrelation function is  $\rho(\tau) = \rho^\tau \cos(\lambda\tau) \left(1 + \frac{1-\rho^2}{1+\rho^2} \tau\right)$ . When  $\lambda=0$ ,  $\cos(\lambda\tau)=1$  for any  $\tau$ . We can compare the autocorrelation functions implied by the two models to that of the observed beef cow inventory series in Table 3. Here we see that the second order stochastic cycle model more closely represents the observed autocorrelation of the beef cow inventory series. In fact for any  $\lambda > 0$ , at some point the implied autocorrelation function will become negative. This feature may not be consistent with the data generating process.

## Conclusion and Discussion

We began our analysis on the dynamics of beef cow numbers with a basic random walk (first-order) stochastic cycle model utilizing semi-annual data of beef cow inventories over a recent 40 year period. Results from the augmented first-order stochastic model imply a cattle cycle of 16.54 years with an asymptotic standard deviation of 1.63 years. We find that a second-order random walk stochastic cycle model provides a better representation of beef cow numbers both in terms

of fit as measured by the log likelihood and implied  $R^2$  and fit as measured by the Ljung-Box Q statistic. However this model yields the unexpected result that the beef cattle cycle disappears.

This raises the question: will there continue to be expansion and contraction of the beef cow herd? Our model does not suggest otherwise. So depending on where a starting point is chosen, there will likely be some pattern of beef cow numbers that allows a cyclical interpretation. The value of our stochastic cycle model is to inform analysts that, using recent data, the cyclical movements are not highly predictable and probably do not conform to a putative nine to twelve-year beef cow cycle. Yes, there are patterns in beef cow numbers that reasoned analysts may interpret as cycles, but our models show there is currently no regularity to cattle cycles such as existed in the past.

For example, the Livestock Marketing Information Center (LMIC) has identified three complete cattle cycles since 1979. Using plots of annual data on total cattle inventory, the LMIC has suggested an eleven year cycle beginning in 1979, a 14 year cycle beginning in 1990, and a ten year cycle beginning in 2004. But unlike earlier patterns in total cattle inventories, the distances from peaks to troughs are muted so as to make cycle identification somewhat subjective. In fact, a basic stochastic cycle analysis of these data over the last 41 years actually identifies an infinitely long cycle even though the dynamic properties of the total annual cattle inventory series differ markedly from those of the semi-annual beef cow inventory series.

Although the identification of commodity cycles can be viewed as purely a statistical exercise, the presence of cycles has important implications. Beveridge and Nelson (1981) have pointed out that the observation of cyclical components in economic time series "has played an important role in shaping our thinking about economic phenomena" (p151.) In this case, we observe either a significant lengthening of the cattle cycle or its complete disappearance. Typically cycles become longer before they disappear. Whereas the popular press has already buried the cattle cycle (Speer, 2014) given the observed patterns in herd size since the 1990's, the second order stochastic cycle model shows its disappearance.

This leads to speculation as to why in recent decades the cycle does not follow the approximate ten year cycle observed over most of the 20th Century. Since 2000 there have been a number of events which have been linked to a lengthening cattle cycle—specifically due to a long trend of decreasing number of beef cows. There were a number of droughts in major cow-calf producing areas from 2000 to 2008 and again in the southern plains from 2010 to 2013 (Petry, 2015). Higher row crop prices and the consequent expansion of crop production may also have led to a reduction in cow-calf operations. According to the 2012 Census of Agriculture, there was a decline of almost 175,000 cow calf operations in the previous 20 years with a bulk of these operations having less than 50 cows. These exogenous shocks hitting the cattle industry could partly explain the change in patterns of beef cow numbers.


While it is debateable as to whether rational expectation theory holds in the cattle industry, our stochastic cycle models identify a lengthening or disappearing cattle cycle; and that certainly gives more credence to a rational expectations paradigm. It will be a matter of time to determine if the cycle re-emerges and reverts to a traditional cycle length of approximately ten years.

*[First submitted February 2020; accepted for publication August 2020.]*


## References

- Aadland, D. (2004). "Cattle Cycles, Heterogenous Expectations and the Age Distribution of Capital." *Journal of Economic Dynamics and Control* 28:1977-2002.  
<https://doi.org/10.1016/j.jedc.2003.08.003>
- Baak, S. J. (1999). "Tests for Bounded Rationality with a Linear Dynamic Model Distorted by Heterogeneous Expectations." *Journal of Economic Dynamics and Control*, 23(9-10), 1517-1543. [https://doi.org/10.1016/S0165-1889\(98\)00082-7](https://doi.org/10.1016/S0165-1889(98)00082-7)

- Beveridge, S. and Nelson, C. R. (1981). "A New Approach to Decomposition of Economic Time Series into Permanent and Transitory Components with Particular Attention to Measurement of the "Business Cycle"." *Journal of Monetary Economics* 7: 151-74.
- Chavas, Jean-Paul. (2000) "On Information and Market Dynamics: The Case of the US BeefMarket." *Journal of Economic Dynamics and Control* 24.5-7 (2000): 833-853. [https://doi.org/10.1016/S0165-1889\(99\)00027-5](https://doi.org/10.1016/S0165-1889(99)00027-5)
- Harvey, A. C. (1985). "Trends and Cycles in Macroeconomic Time Series." *Journal of Business and Economic Statistics* 3: 216-227.
- Harvey, A. C. (1989). *Forecasting, Structural Time Series Models and the Kalman Filter*. New York: Cambridge University Press.
- Harvey, A. C. and Trimbur, T. M. (2003). "General Model-Based Filters for Extracting Cycles and Trends in Economic Time Series." *The Review of Economics and Statistics* 85: 244-255. <https://doi.org/10.1162/003465303765299774>
- Hodrick, R. J. and Prescott, E. C. (1997). "Postwar U.S. Business Cycles: An Empirical Investigation." *Journal of Money, Credit and Banking* 24: 1-16. DOI: 10.2307/2953682
- Howitt, P. and McAfee, R. P. (1992). "Animal Spirits." *The American Economic Review* 82: 493-507.
- Jarvis, L. S. (1974). "Cattle as Capital Goods and Ranchers as Portfolio Managers: An Application to the Argentine Cattle Sector." *Journal of Political Economy* 82: 489-520.
- Nerlove, M. and Fornari, I. (1998) "Quasi-rational Expectations, an alternative to Fully Rational Expectations: An Application to US Beef Cattle Supply." *Journal of Econometrics* 83: 129-161. [https://doi.org/10.1016/S0304-4076\(97\)00067-5](https://doi.org/10.1016/S0304-4076(97)00067-5)
- Parker, P. S. and Shonkwiler, J. S. (2014). "On the Centenary of the German Hog Cycle: New Findings." *European Review of Agricultural Economics* 41: 47-61. <https://doi.org/10.1093/erae/jbt017>
- Petry, T. (2015). "Spotlight on Economics: The Cattle Cycle Revisited." *Agriculture Communication*, North Dakota State University.
- Rosen, S. (1987). "Dynamic Animal Economics." *American Journal of Agricultural Economics* 69: 547-557.
- Rosen, S., Murphy, K. M. and Scheinkman, J. A.(1994). "Cattle Cycles." *Journal of Political Economy* 102: 468-492.
- Rucker, R. R., Burt, O. R., and LaFrance, J. T. (1984). "An Econometric Model of Cattle Inventories." *American Journal of Agricultural Economics* 66: 131-144.
- Slutzky, E. (1972) "The Summation of Random Causes as the Source of Cyclic Processes." *Econometrica* 5: 105-146.
- Speer, N. (2014). "Industry at a Glance: Changes in Beef Cow Inventory vs. Changes in Calf Price." *Beef* June, 24.
- Trimbur, Thomas M. "Properties of Higher Order Stochastic Cycles." *Journal of Time Series Analysis* 27.1 (2006): 1-17. <https://doi.org/10.1111/j.0143-9782.2005.00462.x>
- White, H. (1982). "Maximum Likelihood Estimation of Misspecified Models." *Econometrica* 53: 1-16.



**Figure 1 U.S. Annual Beef Cow Numbers, 1920 – 1981, Corrected for Mean and Trend**



**Figure 2 U.S. Semi-Annual Beef Cow Numbers, 1979 – 2018, Corrected for Mean, Trend, and Seasonality.**

**Table 1. First Order Cycle Estimation Results**

Parameter	First Order Cycle I		First Order Cycle II	
	Estimate	Std.Err. <sup>a</sup>	Estimate	Std.Err.
$\mu_0$	323.800	18.030	337.810	15.338
$\beta$	10.310	5.194	5.872	4.030
$\delta$	-6.750	0.347	-9.828	1.156
$\sigma_\varepsilon$	0		0	
$\lambda$	0.190	0.023	0.190	0.019
$\rho$	0.943	0.015	0.948	0.014
$\sigma_\kappa = \sigma_\nu$	1.677	0.327	1.665	0.298
$\sigma_\eta$	2.568	0.377	2.341	0.347
$\psi_0 = \psi_0^*$	51.203	16.490	43.580	13.528
$\Delta y_{t-3}$			0.234	0.091
Log-likelihood	-206.980		-202.260	
LBQ P-value	0.030		0.258	
R <sup>2</sup>	0.983		0.985	

<sup>a</sup>Robust standard errors

*Note:* The dependent variable is U.S. semi-annual beef cow inventory from 1979 to 2019.  $\mu_0$ ,  $\Psi_0$ , and  $\Psi_0^*$  represent the initial values for  $\mu_t$ ,  $\Psi_t$ , and  $\Psi_t^*$ , respectively.  $\beta$  is the drift term within the random walk component,  $\mu_t$ .  $\delta$  is the coefficient on the semi-annual dummy variable.  $\lambda$  represents cycle frequency, and  $\rho$  is the damping factor of the cycle.  $\sigma_\varepsilon$ ,  $\sigma_\eta$ ,  $\sigma_\kappa$ , and  $\sigma_\nu$  are standard deviations for disturbance terms  $\varepsilon$ ,  $\eta$ ,  $\kappa$ , and  $\nu$ , respectively.  $\Delta y_{t-3}$  is the first difference between  $y_{t-3}$  and  $y_{t-4}$ .

**Table 2. Second Order Cycle Estimation Results**

Parameter	Second Order Cycle I		Second Order Cycle II	
	Estimate	Std.Err. <sup>a</sup>	Estimate	Std.Err.
$\mu_0$	349.260	5.924	358.190	5.218
$\beta$	-5.003	4.086	-7.067	3.377
$\sigma_\varepsilon$	0		0	
$\lambda$	0		0	
$\rho$	0.792	0.041	0.791	0.030
$\sigma_\kappa = \sigma_\kappa$	1.978	0.469	2.119	0.585
$\sigma_\eta$	2.153	0.473	1.715	0.447
$\Psi_0 = \Psi_0^*$	28.445	5.788	24.984	4.618
$\Delta y_{t-3}$			0.217	0.066
Log-likelihood	-204.540		-198.730	
LBQ P-value	0.076		0.934	
R <sup>2</sup>	0.984		0.986	

<sup>a</sup>Robust standard errors

Note: The dependent variable is U.S. annual beef cow inventory from 1979 to 2019.  $\mu_0$ ,  $\Psi_0$ , and  $\Psi_0^*$  represent the initial values for  $\mu_t$ ,  $\Psi_t$ , and  $\Psi_t^*$ , respectively.  $\beta$  is the drift term within the random walk component,  $\mu_t$ .  $\lambda$  represents cycle frequency, and  $\rho$  is the damping factor of the cycle.  $\sigma_\varepsilon$ ,  $\sigma_\eta$ ,  $\sigma_\kappa$ , and  $\sigma_v$  are standard deviations for disturbance terms  $\varepsilon$ ,  $\eta$ ,  $\kappa$ , and  $v$ , respectively.  $\Delta y_{t-3}$  is the first difference between  $y_{t-3}$  and  $y_{t-4}$ .

**Table 3. Autocorrelation functions**

Lag= $\tau$	Observed Cow Numbers	First Order Cycle	Second Order Cycle
1	0.931	0.931	0.974
2	0.906	0.835	0.916
3	0.799	0.718	0.841
4	0.744	0.586	0.758
5	0.623	0.446	0.673
6	0.569	0.304	0.591
7	0.450	0.165	0.514
8	0.395	0.034	0.444
9	0.281	-0.085	0.381
10	0.239	-0.189	0.325